

Geomechanical Aspects of Subsidence in Eastern Maine

Fitzhugh T. Lee and Sharon F. Diehl

*U.S. Geological Survey
Box 25046, MS 966
Denver Federal Center
Denver, Colorado 80225*

ABSTRACT

Information from several sources indicates that the eastern Maine coast is subsiding. Recent releveling surveys indicate a current rate of subsidence in eastern Maine as much as 9 mm per year relative to Bangor. The anomalously rapid rise in sea level accompanying the subsidence has led to shoreline erosion, landsliding, and submergence of archaeological sites and historical manmade structures. In situ stress measurements, well-bore breakouts, and earthquake focal mechanisms show that the maximum principal stress is horizontal, strikes N 80° E, and is correctly oriented to produce thrusting on north-south faults, right-lateral motion with minor thrusting on northeast and east-northeast faults, and major thrusting with left-lateral motion on northwest-striking faults.

Maximum seismicity in Maine occurs in the subsiding zone which also includes the maximum (positive) gravity anomaly in the State. Epicenters are diffusely aligned with the Oak Bay fault which strikes N 20° W. Deflection of isobase lines drawn on ice-marginal deltas forms a trough whose axis strikes approximately parallel to the Oak Bay fault. An acceptable contemporary crustal deformation model should incorporate zones of contrasting rock density and intensity of fracturing, localized seismicity, east-west tectonic compression, and weak compression or extension in the north-south direction.

INTRODUCTION

Crustal subsidence in coastal New England has been recognized from first-order leveling surveys for more than 50 yrs (U.S. Department of Commerce, 1973; Tyler and Ladd, 1980). Geomorphic and cultural evidence suggest that subsidence has occurred over a much longer period, perhaps for 3,000 yrs B.P. (Thompson, 1973; Borns, 1980; Thompson and Kelley, 1983). After subtracting the calculated eustatic sea level rise, Thompson (1973) used radiocarbon dates from basal salt-marsh peat at Addison, Maine, to show a sea level rise of about 4 m in the last 3,000 yrs, with some indication that the rate of rise has been decreasing.

Other methods used to detect crustal movement include submergence of colonial dikes on salt marshes and of harbor facilities, submergence of 3,000- to 5,000-yr-old Indian shell mounds, and changes of altitudes of glacial-marine deltas and beach deposits resulting in displacement of the postglacial-uplift isobases (Anderson and others, 1984). These authors report that

crustal subsidence during historical times was greatest in the Machias-Eastport area. The amount of subsidence has been best established for the last 200 yrs during which time the sea has transgressed over manmade structures in coastal Maine. Most rapid present-day subsidence is located in eastern Washington County where a maximum relative rate of 0.91 m per century has been measured (Tyler and Ladd, 1980). Between 1942 and 1966, subsidence in the area has been as much as 17.5 cm (Tyler and Ladd, 1980).

Less prominent subsidence has been identified in southern York County, Maine (Tyler and Ladd, 1980). Phenomena that may be associated with the subsidence in eastern Maine include localized seismicity, heavily-faulted dense bedrock, and tectonic stresses.

Several economic effects of subsidence have been identified; foremost are continuing coastal erosion and landsliding. In some coastal communities, historic harbor installations are

now under water at high tide. In Eastport, where the subsidence rate is 9 mm per yr, part of the harbor sank as a result of the March 17, 1870 earthquake, and landslides occurred in nearby New Brunswick (Smith and Bridges, 1983).

The purpose of this paper is to summarize structural and tectonic information in eastern Maine and to compare this data with earthquake information, in situ stresses, and gravity interpretations.

REGIONAL GEOLOGY AND TECTONIC HISTORY

Southeastern Maine is underlain by lower and middle Paleozoic sedimentary and volcanic rocks intruded by Devonian plutonic rocks, some of which have batholithic dimensions (Figure 1). There is little granite in the rapidly subsiding area between Machias and Passamaquoddy Bay, a fact which will be discussed in a later section. North and west from the coast in this area the percent of granite bedrock increases significantly (Gates, 1982; Ludman, 1982). Details of stratigraphy, structure, tectonic history and plutonism are given by Gates (this volume). The major faults are shown in Figure 1. Fault strikes in the subsiding zone are shown in Figure 2, and fault distribution is depicted in Figure 3.

Joints and Microfractures

Joints were recorded from exposures of granite at the field test sites (Figure 1), and microfractures were determined in the thin sections prepared from drill hole cores. Joints were designated as major or minor primarily on the basis of continuity and ease of splitting, as recognized in quarries. The attitudes of major joints are similar at all sites: N 85° E, 80° SE; N 80° W, 75° SW; and N 50° W, 82° NE. The joints are widely spaced (1 to 5 m) and typically show little evidence of alteration or movement, although Dale (1907) noted movement along a N 56° E fracture. Faults are not numerous in the Devonian granites in eastern Maine.

Sheeting fractures are encountered in all exposures of the granite. They generally conform to the surface topography and become thicker with depth, ranging from less than 2 cm to more than 2 m at a depth of 17 m.

The metamorphic and volcanic rocks display numerous, closely spaced joints and small faults, usually without apparent pattern. The most common joint sets parallel the regional foliation (N 50°-60° E) and their spacing ranges from 0.2 cm to 0.75 m.

Microfractures occur abundantly in granite from the test sites, and have been analyzed to determine their significance to the interpretation of regional deformation. An analysis of microfractures can provide clues regarding rock anisotropy, deformability, and the possible correlation of micro- and macrostructures (Brace, 1965; Dula, 1981). In some instances, the relative ages of microfractures can be determined. In this study, microfractures include: (1) healed fluid-inclusion planes in

quartz grains, (2) open unlined microfractures occurring as short intragranular cracks to major transgranular features in all minerals, and (3) microfractures that are usually found in feldspar grains and lined with alteration products such as sericite, calcite, or iron oxide. Cleavage planes are often disregarded in a microfracture study, but if there appeared to be a mechanical loss of cohesion along a cleavage surface, especially in the case of a transgranular microfracture paralleling a cleavage plane, its orientation was included in the compilation. Both vertical and horizontal thin sections were prepared, and microfracture frequency and orientation were obtained from two traverses along the length of the thin section. Fabric features were determined using a five-axis universal stage, and then plotted on the lower hemisphere of an equal area net as poles to their respective planes. Data from vertically oriented thin sections were rotated into the horizontal plane and plotted in the composite diagram (Figure 4).

In addition to ubiquitous flat-lying sheeting fractures, the dominant orientations of microfractures are: N 76° E, 70° NW; N 46° E, vertical; and N 86° E, 24° NW. The N 46° E, and N 86° E, strikes are parallel to planes of easy splitting at sites 2 and 3, respectively. There is a marked dominance of steeply dipping N 45° E to east-west striking microfractures (Figure 4). This strike range of fractures is present on several scales (fault zones, joints, microfractures), and in all rock types it is the most prevalent discontinuity observed at the surface in eastern Maine.

GRAVITY INFORMATION

Several gravity studies of subsurface rocks in eastern Maine have been made at various scales (Kane and Bromery, 1968; Kane and others, 1972; Biggi and Hodge, 1982; Hodge and others, 1982).

In their model of the regional gravity field in Maine, Kane and Bromery (1968) noted that the gravity values increase toward the coast in a steplike fashion from -6 mgals in the northwest to +40 mgals in easternmost Maine. Hodge and others (1982) found that the depth of the density contrasts causing the steep northeast-trending gravity gradients is shallow with the basement becoming increasingly shallower toward the coast.

Because gravity anomalies are directly related to the density of the rock below the location of measurement, maps showing Bouguer gravity anomalies are helpful in the evaluation of the thickness of plutons and the distribution of rocks of different densities. Gates (this volume) shows a belt of relatively high-density rocks (positive anomalies) trending northeasterly along the coast, and the eastern part of this belt approximately coincides with the area of most rapid subsidence (Tyler, this volume). According to Kane and Bromery (1968), the average density for rocks in the coastal belt is 3.0 g/cm³, whereas it is 2.65 g/cm³ for the belt to the northwest which contains a larger volume of granitic rocks than the coastal belt.

High gravity values in the northeastern part of the coastal belt correspond to areas underlain by mafic rock, and the sharp




Figure 1. Generalized lithologic map showing major faults in eastern Maine. Modified from Gates (1982) and Ludman (1982).




Figure 2. Strikes of 287 steeply dipping faults measured within 1350 km² area centered on A-A' in Figure 1.




Figure 3. Faults counted within 1350 km² area centered on A-A' in Figure 1. Traverse trends N53°W, perpendicular to subsidence contours of Tyler and Ladd (1980)

gravity gradients at the boundaries of the mafic masses in this area show that these rocks are the principal sources of the gravity highs (Kane and Bromery, 1968). These authors interpret




Figure 4. Lower hemisphere equal area diagram of poles to 438 microfractures measured in thin sections taken from granite core samples from sites 1, 2, and 3. Contoured on 0.4, 1.6, 4.0, 5.2, and 6.4 percent per 1 percent area.

gradient-amplitude relations to mean that contrasts in density (across the boundaries separating blocks of contrasting density) must take place at shallow crustal levels.

Gravity modeling by Hodge and others (1982) indicates that the circular- to elongate-shaped exposures of the Jonesboro, Marshfield, and Meddybemps granitic plutons (sites 2, 3, and 1, Figure 1) join at depth to produce an elongate northeasterly trending batholith. This large granitic mass has an average thickness of approximately 3.5 km (Hodge and others, 1982, Table 3).

EARTHQUAKE ACTIVITY AND FOCAL MECHANISMS

Thirty-nine small earthquakes that occurred from 1976 through 1984 in the Calais-Machias area were reported by Lepage and Johnston (1985). Twenty-three are concentrated in the western Passamaquoddy Bay (Figure 5) and range in magnitude from less than 1.0 to 3.9. Epicenters are more densely clustered in this area than elsewhere on the Maine coast. How-




Figure 5. Earthquakes reported in eastern Maine from January 1, 1976 through December 31, 1984. Earthquakes with magnitudes greater than 2.9 occurred after August 11, 1983. Solid lines represent average maximum and minimum principal horizontal stress vectors. Earthquake data from Lepage and Johnston (1985).

ever, insufficient information is available for these small tremors to calculate reliable focal mechanism parameters.

Focal mechanisms reported for several adequately recorded earthquakes in central Maine, northeastern New York, and southeastern Canada are given in Table 1. Several of the tabulated earthquakes have multiple references. References with the most complete data have been used, which usually includes an interpretation of in situ stresses and type of faulting. The Blue Mountain Lake information is a composite solution of several events ("swarms"). The inferred stress directions indicate an east-west subhorizontal orientation of the maximum principal compression axes for the region. Focal depths range from 2 to 15 km and fault motions are characteristically thrust and reverse (dip of fault plane is greater than 45°), although some have a significant strike-slip component. The principal exceptions to the inferred east-west maximum principal stresses are events 7 and 10 at Otisfield, Maine, and Lake Fairlee, Vermont, where these stresses trend N 30° W and N 35° E, respectively.

The largest earthquakes closest to the subsiding zone (events 2, 4, 6, 8, Table 1) suggest horizontal pressure axes oriented east-west, and would be consistent with thrusting on north-south striking fault surfaces. For event 4, Yang and Aggarwal (1981) determined that the thrust surface strikes N 11° E and dips 48° NW. Ebel and McCaffrey (1984) analyzed the 1983 Dixfield, Maine earthquake (event 6, Table 1) and determined that the main shock showed thrust motion on north-south striking fault surfaces that dip 45° W. From a teleseismic analysis of

event 2, Choy and others (1983) found that the direction of movement was up-dip on a west-dipping, north-northeast striking fault plane. According to these authors, the steep dip (65°) of the inferred fault surface suggests that the earthquake occurred on a preexisting fault that at one time was a normal fault.

These focal mechanisms are consistent with thrust motion on approximately north-south striking moderate to gently dipping surfaces with a horizontal east-northeast to east-west maximum principal stress. A strike-slip component is also present on some fault surfaces. The 1982 Gaza, New Hampshire, earthquake (event 12, Table 1) is exceptional in producing strike-slip motion on a N 20° E - 80° SE oriented fault surface.

Considering the diversity of fault attitudes, ages of faulting, and variety of rock types in northeastern New England and adjacent Canada, the consistency of pressure axes of focal mechanism solutions is good.

IN SITU STRESSES

In situ measurements of rock stress were made in the subsiding zone at the three sites shown in Figure 1. Because of the intensity of fracturing and scarcity of granitic rocks in the most rapidly subsiding zone, a suitable site could not be located there. If all three test sites are located in the same northeasterly elongated, partially buried granite pluton, then regional significance may be attributed to a discussion of in situ stresses and contemporary rock deformation. The composition of the rocks at the

TABLE 1. EPICENTER AND SOURCE DATA FOR EARTHQUAKES IN NORTHERN NEW ENGLAND AND SOUTHEASTERN CANADA (Leaders (---) indicate data missing; R=reverse; T=thrust; SS=strike slip; H=horizontal)

Event No.	Geographic region	Date	Lat. (°N)	Long. (°W)	Depth (km)	Magnitude	Maximum principal stress axis		Minimum principal stress axis		Dist. from Machias, ME (km)	Type of fault movement	References
							Azimuth	Plunge	Azimuth	Plunge			
1	La Malbaie, Quebec	6/30/74	47.7	69.8	15	*2.0	N86°W	5°N	N4°E	83°S	390	R	Leblanc and Buchbinder (1977)
2	Miramichi, New Brunswick	1/9/82	47.0	66.6	6-10	5.7	E-W	H	N-S	70°S	250	R/SS	Adams and Wetmiller (1983); Hasegawa (1983); Choy and others (1983)
3	Maniwaki, Quebec	7/12/75	46.3	76.1	17-19	4.3	N50°E	19°N	N40°W	---	695	R/SS	Horner and others (1975)
4	Maine-Quebec	6/15/73	45.28	70.97	7	4.8	N77°E	H	N13°W	90°	286	T	Yang and Aggarwal (1981)
5	Altona, NY	6/9/75	44.9	73.67	10	4.2	N73°E	8°N	N17°W	84°S	518	R	Aggarwal and others (1977)
6	Dixfield, ME	5/29/83	44.51	70.41	1.8-2.4	4.4	E-W	H	N-S	90°	235	T	Ebel and McCaffrey (1984)
7	Otisfield, ME	1/4/78	44.04	70.51	3	3.2	N30°W	30°N	N70°W	70°S	256	R	Graham and Chiburis (1980)
8	Bath, ME	4/18/79	43.85	69.76	3	4.0	E-W	H	N-S	90°	199	T	Pulli and Toksoz (1981)
9	Blue Mountain Lake, NY	5/7/71	43.88	74.33	2-3	3.1-3.6	N71°E	18°N	N19°W	73°S	515	T	Sbar and Sykes (1977)
10	Lake Fairlee, VT	5/5/77	43.86	72.26	6	2.1	N35°E	30°N	N55°W	35°S	398	R/SS	Graham and Chiburis (1980)
11	Boonville, NY	6/6/80	43.60	75.10	2	3.5	N75°E	10°N	N42°E	80°S	650	T	Pulli and Toksoz (1981)
12	Gaza, NH	1/19/82	43.52	71.61	3	4.6	N65°E	H	N25°W	90°	357	SS	Pulli, J. J. (written commun., 1984)
13	Hopkinton, NH	12/25/77	43.19	71.65		3.2	N76°W	15°N	N-S	35°S	381	T/SS	Graham and Chiburis (1980)
14	Candia, NH	4/23/79	43.04	71.24	2	3.1	E-W	H	N-S	90°	350	T/SS	Pulli and Toksoz (1981)
Average							N83°E	10°N	N14°W	74°S			

*Largest of 35 events over a 7-week period.




Figure 6. Modal analyses of granite from three field sites. Samples from Meddybemps Lake (site 1) contain some admixing of more basic country rock.

test sites is shown in Figure 6. The slightly more mafic composition of the granite at site 1 is caused by local mixing of gabbroic country rock (Amos, 1963). On the basis of similarity of the petrographic characteristics, this author determined that the granites belong to the same plutonic complex and were probably emplaced during a relatively short time interval. This conclusion agrees with the similar 403-407 m.y. age determinations of four granites in southeastern Maine (Faul, 1960).

The objectives of the stress-measurement program were as follows:

1. Determine directions and magnitudes of principal horizontal stresses.
2. Measure vertical rock stresses and compare results with vertical gravitational stresses.
3. Compare shallow measurements of horizontal rock stresses with stress directions inferred from earthquake focal mechanisms.
4. Suggest potential subsidence mechanisms.

Results of Stress Measurements

Several factors affect shallow rock stresses, the most important of which are temperature, topography, rock fabric, and geometric stress concentrations from quarry walls and corners. These factors were carefully considered in site selection, and drill holes were located in areas of subdued topography on bedrock surfaces distant from rock walls, faults, and fracture zones. The most severe diurnal and annual temperature changes occur in the upper 3 to 4 m of a rock mass (Hooker and Duvall, 1971). Therefore, the deepest measurements at the three sites

(average of 5.03 m) are regarded as the most reliable. No rock mass is devoid of fractures; hence, the influence of faults, joints, and microfractures on strain relief should be examined. Except at site 1, joints are so widely spaced that they were not encountered in the boreholes and should have a negligible effect on the measurements. The joints were commonly tightly closed with no alteration. The influence of microfractures on horizontal strain-relief measurements could cause directional differences in Young's modulus (stiffness). The ratio of maximum to minimum stiffness in the horizontal plane for the deepest measurement at each site is: site 1, 1.16; site 2, 1.15; and site 3, 1.11. Except at site 2, the direction of maximum stiffness coincides with the major microfracture directions (Figure 4; Table 2). At site 2, the only identified structural fabric element, which is close to the maximum stiffness direction, is a 2 m-thick aplite dike which strikes N 5° E, and dips 75° NW. The stiffness anisotropy (average ratio of 1.14 for the deepest measurement) is overwhelmed by the ratio of maximum to minimum horizontal stresses (average ratio of 7.4). Therefore, the strongly deviatoric horizontal stresses, as reported, are little affected by stiffness variations but mainly controlled by the far-field regional stresses.

Horizontal stresses were determined at all three sites by the stress-relief overcoring method using the U.S. Bureau of Mines borehole gage in vertical holes (Hooker and Bickel, 1974). In addition, vertical stresses were measured in a horizontal borehole at site 2. Three horizontal displacement measurements were made in vertical holes at each site at depths ranging from 0.95 to 6.4 m. As expected, Young's modulus values are lowest at site 1 and highest at site 2, where the rock is most massive (Table 2).

Maximum principal horizontal stress directions range from N 56° E, to N 85° W, and average N 79° E. The minimum principal horizontal stress directions range from N 5° E, to N 34° W, and average N 11° W. Stress magnitudes were greatest at site 2 in the Jonesboro Granite where a maximum value of 14.01 MPa was obtained at a depth of 6.4 m. Lowest maximum principal stress magnitudes were found at site 1, where three measurements averaged only 1.39 MPa. At this site, tensile stresses were measured at shallow depths (Table 2). The horizontal stress deviator, or difference in the horizontal principal stresses, is shown in Table 2. Significant stress deviation may be an indication of potential instability along fracture surfaces due to the presence of shear stresses.

The vertical stress was measured in a wall of the quarry at site 2 in a subhorizontal hole (+3°) at a distance of 4 m from the collar. The vertical rock column above the measurement position is approximately 6 m, which corresponds to a vertical gravitational stress of 0.14 MPa. This is in good agreement with the measured value of 0.11 MPa. This measurement suggests that near the earth's surface, the minor principal stress (least compression) is vertical and the maximum and intermediate principal stresses are horizontal.

The approximate east-west maximum horizontal compression that was calculated from our stress-relief overcoring meas-

TABLE 2. IN SITU STRESSES MEASURED IN THE SUBSIDENCE ZONE

Site No. $\sigma_1 + \sigma_2$ 2	Depth (m)	Maximum principal horizontal stress (σ_1)		Minimum principal horizontal stress (σ_2)		Mean horizontal stress (MPa)	Stress deviator ($\sigma_1 - \sigma_2$)	Average Young's modulus (GPa)	Direction of maximum Young's modulus	Rock type
		(MPa)	(Strike)	(MPa)	(Strike)					
1	3.9	0.08	N 78° E	-3.82	N 12° W	-1.87	3.92	27.9	N 80° E	Granite
	4.3	-0.12	N 84° E	-2.11	N 6° W	-1.12	1.99	33.6	N 70° W	Gneissic granite
	5.2	4.20	N 66° E	0.29	N 24° W	2.25	3.91	30.8	N 80° E	Gabbro
Average 2	4.5	1.39	N 76° E	-1.88	N 14° W	-0.25	3.27	30.8	E - W	
	3.45	12.03	N 85° E	3.25	N 5° W	7.64	8.87	51.2	N 10° W	Granite
	4.19	10.95	N 87° E	2.53	N 2° W	6.74	8.42	50.7	N 10° W	Granite
	6.45	14.01	N 85° W	2.45	N 5° E	8.23	11.56	50.1	N 10° W	Granite
Average	4.7	12.33	N 89° E	2.74	N 1° W	7.54	9.59	50.7	N 10° W	
3	0.95	11.67	N 56° E	5.15	N 34° W	8.41	6.52	29.9	N 70° E	Granite
	2.97	8.96	N 82° E	3.43	N 8° W	6.20	5.53	34.3	N 70° E	Granite
	3.43	10.26	N 78° E	5.25	N 12° W	7.76	5.01	34.7	N 80° W	Granite
Average	2.45	10.30	N 72° E	4.61	N 18° W	7.46	5.69	32.97	N 80° E	
Average for sites 1-3	3.87	8.01	N 79° E	1.82	N 11° W	4.92	6.19	38.13		

Note: Site locations are shown on Figure 1. Negative values denote tensile stresses.

urements is in good agreement with the orientation of tectonic forces estimated by other methods. An east-west horizontal principal compression is also shown from an analysis of well bore breakouts (sidewall spalling in near-vertical holes) in six deep wells in the Georges Bank (Zoback and others, in press; R. L. Dart, written commun., 1985). Further evidence of east-west compression was obtained by Tyler and Leick (1985; this volume) from a geodetic survey in the Calais area. These authors obtained first order accuracy using macrometer-satellite distance measurements. Their comparison of 1983 values with 1890 triangulation survey data indicates a strong possibility of significant crustal motion although the data are not sufficient to provide unequivocal evidence of crustal deformation.

The stress-resolution procedure of Swolfs (1984) is helpful in estimating the sense of potential fault movement. This procedure utilizes a triangular diagram on which the three principal stresses are represented by a single point that designates the preferred mode and orientation of faulting (Figure 7). Pore-water pressures in the massive, practically impermeable granites are considered to be low so that the total stresses (S_1 , S_2 , and S_3) are approximately equal to the effective stresses (σ_1 , σ_2 , and σ_3). The measured stresses plot in the thrust faulting section of the diagram ($S_y S_x S_z$). If the vertical stress (S_z) increases with depth relative to the minimum horizontal stress (S_x), as has been observed at several locations in the depth range <1km to 5 km, a component of strike-slip motion will be present on appropriately oriented faults (Swolfs, 1984). This is shown by the dashed line on Figure 7. A component of strike-slip motion was reported for several of the events listed in Table 1.




Figure 7. Triangular stress diagram showing structural definition of six panels in terms of the preferred mode and orientation of faulting (after Swolfs, 1984). The average stresses from Table 2 are plotted in percent along the appropriate total stress sides ($S_y S_x S_z$). A thrust-faulting regime is indicated. Dashed line is explained in text.

DISCUSSION AND CONCLUSIONS

The nature and origin of tectonic forces along continental margins has been inferred from geophysical and geomorphic data. For example, Yang and Aggarwal (1981) found that the parallelism between the maximum principal horizontal stress, determined from focal plane mechanisms, and the age gradient of oceanic lithosphere spreading, was parallel to the gradient of offshore magnetic anomalies determined by Pitman and others (1974). Intraplate movements near the Atlantic margin of the northeastern United States may be generated by an east-west horizontal compression reflecting the gravitational drive acting at the continental margin, owing to horizontal density variations in the oceanic lithosphere as it cools and moves away from spreading centers (Yang and Aggarwal, 1981). The stress generated by this gravitational force was modeled by these authors to be in the direction of absolute plate motion, of less than a few hundred bars magnitude (<20 MPa) and could be readily modified by local effects. Heller and others (1982) stated that the regional tectonic regime along the United States Atlantic margin has been consistent since rifting (mid-Mesozoic), but they could not identify the driving force of episodic Cenozoic subsidence.

Crustal loading from dense, thick volcanic rocks, fracturing, and the orientation of tectonic stress directions relative to Acadian or Taconic faults are factors that could facilitate subsidence in eastern Maine. The decrease of subsidence to the northwest inland from the coast coincides with the decreases in rock density and severity of fracturing in the same direction. At shallow depths, the contemporary maximum horizontal stress (N 79° E) is correctly oriented to produce "pure" thrust on north-south faults, right-lateral motion with minor thrusting on north-east and east-northeast faults, and left-lateral motion with major thrusting on northwest-striking faults. Seismicity is a maximum in Maine in the area of maximum subsidence, and epicenters also show a diffuse pattern along the Oak Bay fault (Figure 1). The abrupt change in elevations of glaciomarine ice-marginal deltas and wave-cut terraces in eastern Maine (Thompson and others, this volume) has produced a deflection of isobases which also suggests that subsidence may be controlled by north and northwesterly striking faults. Alternatively, this deformation could be accounted for by sharp flexing. In the vicinity of the Oak Bay fault, the isobases delineate a trough whose axis is approximately parallel to the fault. Contemporary subsidence contours in New Brunswick and Nova Scotia trend north to northwest (Lambert and Vanicek, 1979), which suggests that northerly striking structures may be involved.

Neotectonic regional crustal behavior in eastern Maine and adjacent Canada is not sufficiently well understood to quantify the role of individual structural elements or blocks in the current subsidence. An acceptable qualitative model for crustal subsidence in eastern Maine should incorporate zones of contrasting rock density, intensity of fracturing, and localized seismicity in addition to tectonic compression directed east-west, and weak

compression or extension in the north-south direction. It is clear that future leveling should reconcile present differences in crustal-velocity contour magnitude and trend across the national boundary. Additionally, future research should include establishing a portable microseismic array in the area of most rapid subsidence in order to evaluate more accurately local fault activity at shallow depths.

ACKNOWLEDGMENTS

The writer is thankful for the help of several individuals in expediting the field and office investigations. Allan Ludman and Olcott Gates spent time in the field explaining the geology of the Fredericton and Eastport 2° quadrangles and suggesting drill sites. David Stevens of the H. E. Fletcher Company made available drill sites in the Hall's Ridge quarry at Jonesboro, Maine, and aided us in our operations. John E. Ebel, Henri S. Swolfs, and R. Ernest Anderson reviewed the first draft of the manuscript and suggested several improvements. The Maine Geological Survey supplied field vehicles and logistical support, and the U.S. Nuclear Regulatory Commission provided partial funding for the project. Danny R. Miller and C. Bruce Boydston obtained in situ strain-relief measurements. Danny R. Miller assisted with office calculations, drafting, and accomplished numerous logistical chores.

REFERENCES CITED

- Adams, J., and Wetmiller, R. J., 1983, Conjugate thrust faulting during the Miramichi, New Brunswick earthquake sequence of 1982 -- Its geometry, geological control, surface expression and mechanism (abs.): *Earthquake Notes*, v. 54, p. 84.
- Aggarwal, Y. P., Yang, J. P., and Cranswick, E., 1977, Seismological investigation in the Adirondacks and environs: *Geological Society of America Abstracts with Programs*, v. 9, p. 234.
- Amos, D. H., 1963, Petrology and age of plutonic rocks, extreme southeastern Maine: *Geological Society of America Bulletin*, v. 74, p. 169-194.
- Anderson, W. A., Kelley, J. T., Thompson, W. B., Borns, H. W., Jr., Sanger, D., Smith, D. C., Tyler, D. A., Anderson, R. S., Bridges, A. E., Crossen, K. J., Ladd, J. W., Andersen, B. G., and Lee, F. T., 1984, Crustal warping in coastal Maine: *Geology*, v. 12, p. 677-680.
- Ballard, R. D., and Uchupi, E., 1975, Triassic rift structure in the Gulf of Maine: *American Association of Petroleum Geologists Bulletin*, v. 59, p. 1041-1072.
- Biggi, R. J., and Hodge, D. S., 1982, Gravity studies over the Bays of Maine igneous complex, in Hussey, A. M., ed., *Maine Geology: Geological Society of Maine, Bulletin 2*, p. 1-15.
- Borns, H. W., Jr., 1980, Summary of research activity in crustal warping study, in Thompson, W. B., ed., *New England Seismotectonic Study activities in Maine during fiscal year 1980: Maine Geological Survey Report to U. S. Nuclear Regulatory Commission*, p. 75-78.
- Brace, W. F., 1965, Relation of elastic properties of rocks to fabric: *Journal of Geophysical Research*, v. 70, p. 5657-5667.
- Choy, G. L., Boatwright, J., Dewey, J. W., and Sipkin, S. A., 1983, A teleseismic analysis of the New Brunswick earthquake of January 9, 1982: *Journal of Geophysical Research*, v. 88, p. 2199-2212.
- Dale, T. N., 1907, The granites of Maine: *U.S. Geological Survey, Bulletin 313*, 202 p.

- Dula, W. F., Jr., 1981, Correlation between deformation lamellae, microfractures, macrofractures, and in situ stress measurements, White River uplift, Colorado: *Geological Society of America Bulletin*, v. 92, p. 37-46.
- Ebel, J. E., and McCaffrey, S. J., 1984, Hypocentral parameters and focal mechanisms of the 1983 earthquake near Dixfield, Maine: *Earthquake Notes*, v. 55, p. 21-24.
- Faul, H., 1960, Geologic time scale: *Geological Society of America Bulletin*, v. 71, p. 637-644.
- Gates, O., 1982, Brittle fractures in the Eastport 2-degree sheet, Maine: Augusta, Maine, Maine Geological Survey Open-File Report 82-29, 15 p.
- Graham, T., and Chiburis, E. F., 1980, Fault plane solutions and the state of stress in New England: *Earthquake Notes*, v. 51, p. 3-12.
- Hasegawa, H. S., 1983, Surface wave analysis of the magnitude 5.7 Miramichi, New Brunswick, earthquake of 09 January 1982 (abs.): *Earthquake Notes*, v. 54, p. 84.
- Heller, P. L., Wentworth, C. M., and Poag, C. W., 1982, Episodic post-rift subsidence of the United States Atlantic continental margin: *Geological Society of America Bulletin*, v. 93, p. 379-390.
- Hodge, D. S., Abbey, D. A., Harbin, M. A., Patterson, J. L., Ring, M. J., and Sweeney, J. F., 1982, Gravity studies of subsurface mass distributions of granitic rocks in Maine and New Hampshire: *American Journal of Science*, v. 282, p. 1289-1324.
- Hooker, V. E., and Bickel, D. L., 1974, Overcoring equipment and techniques used in rock stress determination: U.S. Bureau of Mines Information Circular 8618, 32 p.
- Hooker, V. E., and Duvall, W. I., 1971, In situ rock-temperature: U.S. Bureau of Mines Report of Investigations RI 7589, 12 p.
- Horner, R. B., Stevens, A. E., Hasegawa, H. S., and Leblanc, G., 1975, The Maniwaki, Quebec, earthquake of July 12, 1975 (abs.): *Earthquake Notes*, v. 46, p. 48.
- Kane, M. F., and Bromery, R. W., 1968, Gravity anomalies in Maine, in Zen, E., White, W. S., Hadley, J. B., and Thompson, J. B., Jr., eds., *Studies of Appalachian geology, northern and maritime*: New York, Interscience Publishers, p. 415-423.
- Kane, M. F., Yellin, M. J., Bell, K. G., and Zietz, I., 1972, Gravity and magnetic evidence of lithology and structure in the Gulf of Maine region: U.S. Geological Survey, Professional Paper 726-B, 22 p.
- Lambert, A., and Vanicek, P., 1979, Contemporary crustal movements in Canada: *Canadian Journal of Earth Sciences*, v. 16, p. 647-668.
- Leblanc, G., and Buchbinder, G., 1977, Second microearthquake survey of the St. Lawrence Valley near La Malbaie, Quebec: *Canadian Journal of Earth Sciences*, v. 14, p. 2778-2789.
- Lepage, C. A., and Johnston, R. A., 1985, Earthquakes in Maine -- October 1975 to December 1984: Augusta, Maine, Maine Geological Survey Open-File Map 85-71, scale 1:500,000.
- Ludman, A., 1981, Significance of transcurrent faulting in eastern Maine and location of the suture between Avalonia and North America: *American Journal of Science*, v. 281, p. 463-483.
- Ludman, A., 1982, Bedrock geology of the Fredericton 2-degree Quadrangle, Maine: Augusta, Maine, Maine Geological Survey, Open-File Report 82-30, 16 p.
- Pitman, W. C., III, Larson, R. L., and Herron, E. M., 1974, Map of magnetic lineations of the oceans: New York, Lamont-Doherty Geological Observatory of Columbia University.
- Pulli, J. J., and Toksoz, M. N., 1981, Fault plane solutions for northeastern United States: *Seismological Society of America Bulletin*, v. 71, p. 1875-1882.
- Sbar, M. L., and Sykes, L. B., 1977, Seismicity and lithospheric stress in New York and adjacent areas: *Journal of Geophysical Research*, v. 82, p. 5771-5786.
- Smith, D. C., and Bridges, A. E., 1983, Historical dating of salt marsh dikes in coastal Maine/Historically recorded earthquakes in eastern Maine: Augusta, Maine, Maine Geological Survey, Open-File Report 83-10, 12 p.
- Swolfs, H. S., 1984, The triangular stress diagram -- A graphical representation of crustal stress measurements: U.S. Geological Survey Professional Paper 1291, 19 p.
- Thompson, S. N., 1973, Sea-level rise along the Maine coast during the last 3,000 years: Orono, Maine, University of Maine, M.S. thesis, 65 p.
- Thompson, W. B., and Kelley, J. T., eds, 1983, *New England Seismotectonic Study activities in Maine during fiscal year 1982*: Maine Geological Survey Report to U. S. Nuclear Regulatory Commission, 171 p.
- Tyler, D. A., and Ladd, J. W., 1980, Vertical crustal movement in Maine: Augusta, Maine, Maine Geological Survey, Open-File Report 80-34, 53 p.
- Tyler, D. A., and Leick, A., 1985, St. Croix region crustal strain study: Augusta, Maine, Maine Geological Survey, Open-File Report 85-76, 22 p.
- U.S. Department of Commerce, 1973, Reports on geodetic measurements of crustal movement, 1906-71: National Oceanic and Atmospheric Administration, National Ocean Survey, and National Geodetic Survey, p. 65.
- Yang, J. P., and Aggarwal, Y. P., 1981, Seismotectonics of northeastern United States and adjacent Canada: *Journal of Geophysical Research*, v. 86, no. B6, p. 4981-4998.
- Zoback, M. D., Moos, D., Mastin, J., and Anderson, R. N. (in press), Wellbore breakouts and in situ stress: *Journal of Geophysical Research*, 28 p.