



RECONNAISSANCE  
SURFICIAL GEOLOGY  
OF THE  
**NORTH EAST CARRY  
QUADRANGLE, MAINE**

OPEN-FILE NO. 76-2

BY  
**THOMAS BREWER  
DABNEY W. CALDWELL**

Maine Geological Survey  
DEPARTMENT OF CONSERVATION  
Augusta, Maine 04333  
Walter A. Anderson, State Geologist

1976




Financial assistance provided by the  
Land Use Regulation Commission,  
Department of Conservation.


For more detailed information refer  
to the "Surficial Geology Hand-  
book for Coastal Maine" by W.B.  
Thompson, 1979.

**CORRELATION OF MAP UNITS**


This correlation chart shows the general age  
relationships of surficial deposits. There may be  
considerable overlap in the ages of certain  
deposits in any particular area.




11,000 years ago. Glacier had dis-  
appeared from central and southern  
Maine. Uplift of land had caused  
sea to retreat.



13,000 years ago. Glacier was  
receding rapidly and much of  
southern Maine was ice-free. Land  
was still depressed from weight of  
ice, resulting in extensive marine  
submergence of lowland areas.




13,500 years ago. Continental glacier  
covered most of Maine, but was  
receding from the coastal lowland.  
Sea was in contact with ice margin.



- BEM — Buried end moraine
- BR — Bedrock ridge
- D — Delta
- DR — Drumlins
- DS — Distributary stream
- E — Esker
- EM — End moraine
- IB — Ice block
- K — Kettle
- M — Marine sediments
- ML — Marine limit
- OP — Outwash plain
- S — Seawater
- T — Till

MARINE LIMIT: Shows region of  
Maine covered by sea water during  
period of maximum inundation  
about 13,000 years ago.

Text by W.B. Thompson. Graphics by R.D. Tucker.



| SYMBOL | GEOLOGIC UNIT   | MATERIALS<br>(Listed in decreasing order of abundance)  | TOPOGRAPHY   | ORIGIN   |
|--------|---|---|--|--|
| Qal    | Stream alluvium (flood-plain and stream terrace deposits) | Sand, gravel, and silt. Low to high permeability. Poor to good drainage. Permeability and drainage generally are better in stream terrace deposits than in modern flood-plain sediments.  | Flat to gently sloping.  | Deposited on flood plains and stream beds by postglacial streams. Unit may also include minor alluvial fan deposits at mouths of valleys.  |
| Qs     | Swamp and tidal-marsh deposits                            | Peat, silt, clay, and sand. Poor drainage.  | Flat.  | Formed by accumulation of sediments and organic material in depressions and other poorly drained areas.  |
| Qbd    | Beach and dune deposits                                   | Sand and gravel. High permeability. Materials are well drained, but water table is close to surface.  | Low ridges and mounds, or sloping surface.   | Occurs along modern ocean and lake shores. Includes beach sediments formed by wave and current action, and windblown sand derived from these deposits.   |
| Qe     | Eolian deposits   | Sand. Moderate to high permeability. Good drainage.   | Dune ridges and mounds, or blanket deposit that conforms to surface of underlying unit.  | Windblown sand. Derived from wind erosion of glacial sediments and deposited in late-glacial to postglacial time.  |
| Qta    | Talus deposits  | Large, angular rock fragments.  | Steeply sloping rock piles at the bottoms of cliffs.   | Formed by the accumulation of rock fragments that break loose from a cliff and fall to the slope below.  |
| Ql     | Glacial-lake bottom deposits                              | Silt, clay, and sand, commonly as thin, interstratified layers of various particle sizes. Low to moderate permeability. Poor to fair drainage.  | Flat to gently sloping except where dissected by modern streams.   | Composed largely of sediments that washed out of glacial ice and accumulated on the floors of glacial lakes. Map unit may also include a few non-glacial lake bottom deposits.                 |
| Qbl    | Glacial-lake beach deposits                               | Gravel and sand. Typically thin and of limited extent. High permeability. Good drainage.  | Low ridges or sloping surface. May be associated with wave-cut benches on hillsides.   | Formed by wave erosion of till or other materials along shores of glacial lakes. Lakes have since lowered or drained.  |
| Qps    | Glacial-marine deposits (Presumpscot Formation)           | Silt, clay, and sand. Commonly a clayey silt, but sand is very abundant at the surface in some places. Locally fossiliferous. Map unit includes small areas of till, sand, and gravel that are not completely covered by marine sediment.   | Flat to gently sloping except where dissected by modern streams. Commonly has a branching network of steep-walled stream gullies.  | Composed of sediments that washed out of the Late Wisconsinan glacier and accumulated on the ocean floor. Formed during late-glacial time, when relative sea level was higher than at present. |
| Qop    |   |   |  |  |
| Qbm    | Emerged marine-beach deposits                             | Gravel and sand. High permeability. Good drainage. Typically thin and of limited extent.  | Low ridges or sloping surface. May be associated with wave-cut benches on hillsides.   | Formed by wave erosion of till or other materials during late-glacial time, when relative sea level was higher than at present.  |
| Qgo    | Glacial-stream deposits                                   | Sand and gravel. May include minor till. Commonly overlapped or entirely buried by glacial-marine deposits in the coastal lowland.  | Qgo: Flat (outwash deposits).<br>Qg: Flat-topped landforms, bounded in part by steep sides (kame terraces and kame deltas); hummocky terrain with hills (kames) and depressions (kettles), or ridges (eskers). | Deposited by meltwater streams and currents during melting of the Late Wisconsinan glacier.  |
| Qg     |   | Qgo: High permeability. Good drainage.<br>Qg: Moderate to high permeability. Good drainage.   | Letters indicate good examples of particular landforms:<br>d delta<br>k kame<br>kt kame terrace  | Qgo: Outwash plains and outwash deltas. Deposited beyond the ice margin.<br>Qg: Kames, eskers, and marine or glacial-lake deltas—all formed adjacent to glacial ice (ice-contact deposits).    |
| Qmg    | End-moraine deposits                                      | Till and/or sand and gravel. Commonly overlain by glacial-marine deposits in coastal areas. Permeability and drainage are highly variable, even over short distances in a single moraine.   | Ridges. Commonly arcuate, discontinuous, and in groups. Size ranges: 3.75 ft. high, 20-1,000 ft. wide, and 100 ft. to several miles long.  | Deposited in marginal zone of the Late Wisconsinan glacier, by glacial ice (till) and/or meltwater streams emerging from the ice (sand and gravel).  |
| Qmh    |   | Qmg: Composed mostly of sand and gravel.<br>Qmh: Composed of till, sand, and gravel.<br>Qmt: Composed mostly of till.<br>Qm: Composition not specified.   | Qmh: Hummocky topography.  |  |
| Qm     |   |   |  |  |
| Qt     | Till  | Heterogeneous mixture of sand, silt, clay, and stones. Stratification is rare. Includes two varieties: basal till and ablation till. Basal till is fine grained and very compact, with low permeability and poor drainage. Ablation till is loose, sandy, and stony, with moderate permeability and fair to good drainage. Unit generally overlies bedrock, but may overlie or include sand and gravel. | Commonly a blanket deposit that conforms to bedrock surface. Also forms glacially stream-lined hills (including drumlins), where till thickness locally exceeds 100 feet.                                      | Deposited directly by glacial ice.   |
| rk     | Bedrock outcrops  | Dots show locations of individual outcrops. Ruled pattern indicates areas of many outcrops and/or thin surficial deposits (generally less than 10 ft. thick). Symbol "rk" indicates areas of barren ledge. Outcrops mapped largely by interpretation of aerial photography in off-road areas.   |  |  |
| af     | Artificial fill   | Composed of till, sand and gravel, rock or various man-made materials (mainly trash in large dumps and landfills).  |  |  |
| —      | Contact   | Boundary between adjacent map units. Dashed where inferred.   |  |  |
| —      | Scarp   | Separates stream terrace from modern flood plain and adjacent terraces from each other. Machures on downslope side.   |  |  |
| —      | Drumlin   | Glacially streamlined hill that has been elongated in direction of ice movement. Symbol shows direction of long axis. Generally composed of till and/or bedrock (rarely sand and gravel). Till is very thick in parts of many drumlins.   |  |  |
| →      | Glacial striation locality                                | Point of observation at tip of arrow. Arrow indicates ice-movement direction as inferred from striations (scratches on bedrock caused by glacial abrasion).   |  |  |
| →      | Crescentic mark locality                                  | Point of observation at tip of arrow. Arrow indicates ice-movement direction as inferred from crescentic marks on bedrock surface.  |  |  |
| —      | Grooved till surface                                      | Symbols show lengths and directions of narrow ridges carved in till by flow of glacial ice.   |  |  |
| —      | Area of many large boulders                               |   |  |  |
| —      | Cirque  | A steep-walled, half-bowl shaped basin. Formed by glacial erosion in high mountainous areas of Maine.   |  |  |
| —      | End moraine   | Ridge of till or sand and gravel deposited at margin of glacier. Barbs point in direction of ice movement. Symbol is used in part to indicate moraines that are mostly buried by water-laid glacial sediment, as well as moraines that are too narrow to be outlined by a contact line at the scale of the map.   |  |  |
| 290    | Dip direction of delta forest beds                        | Number is approximate altitude in feet of contact between topset and forest beds, which marks former position of sea level or glacial lake level (generally sea level in coastal Maine). Point of observation at tip of arrow.  |  |  |
| →      | Dip direction of cross bedding in glacial-stream deposits | Indicates direction of flow of glacial meltwater streams.   |  |  |
| >>>    | Crest of esker  | Shows trend of sand and gravel ridge that was deposited in meltwater tunnel beneath glacier. Chevrons point in direction of meltwater flow.   |  |  |
| —      | Kettle  | Depression created by melting of large mass of buried glacial ice and collapse of overlying sediments.  |  |  |
| —      | Meltwater channel   | Channel eroded by glacial meltwater stream. Arrow indicates known or probable direction of stream flow.   |  |  |
| —      | Meltwater channel   | Flow direction not specified.   |  |  |
| —      | Till or sand and gravel pit                               | Letter symbols indicate materials exposed in pit:<br>t till<br>t <sub>l</sub> loose, sandy ablation till<br>t <sub>c</sub> compact, fine-grained basal till<br>f flowtill<br>cy clay<br>st silt<br>s sand<br>ps pebbly to cobbly sand<br>p pebble gravel<br>c cobble gravel<br>b boulder gravel<br>g gravel, undifferentiated<br>rk bedrock<br>rs rottenstone (decomposed bedrock)                      |  |  |
| —      | Bedrock quarry  | Superposed symbols indicate superposition of materials. Read hyphen as "to."<br>X active or unchecked<br>X inactive   |  |  |
| ●      | Source of dated sample                                    | SI-3041<br>Symbols show collection site and laboratory sample number of <sup>14</sup> C dated shells, wood, or other organic material.  |  |  |