

Surficial Geology

Camden Quadrangle, Maine

Surficial geologic mapping by
Woodrow B. Thompson

Digital cartography by:
Robert A. Johnston

Robert G. Marvinney
State Geologist

Cartographic design and editing by:
Robert D. Tucker

Funding for the preparation of this map was provided in part by the U.S. Geological Survey STATEMAP Program, Cooperative Agreement No. G09AC00178.



Maine Geological Survey

Address: 22 State House Station, Augusta, Maine 04333
Telephone: 207-287-2801 E-mail: mgs@maine.gov
Home page: http://www.maine.gov/dor/nrmc/nrmc.htm

Open-File No. 10-6
2010

SURFICIAL GEOLOGY OF MAINE

Continental glaciers like the ice sheet now covering Antarctica probably extended across Maine several times during the Pleistocene Epoch, between about 1.5 million and 10,000 years ago. The slow-moving ice superficially changed the landscape as it scraped over mountains and valleys, eroding and transporting boulders and other rock debris for miles. The sediments that cover much of Maine are largely the product of glaciation. Glacial ice deposited some of these materials, while others washed into the sea or accumulated in meltwater streams and lakes as the ice receded. Earlier stream patterns were disrupted, creating hundreds of ponds and lakes across the state. The map at left shows the pattern of glacial sediments in the Camden quadrangle.


The most recent "Ice Age" in Maine began about 30,000 years ago when an ice sheet spread southward over New England (Stone and Borns, 1986). During its peak, the ice was several thousand feet thick and covered the highest mountains in the state. The weight of this huge glacier actually caused the land surface to sink hundreds of feet. Rock debris frozen into the base of the glacier abraded the bedrock surface over which the ice flowed. The grooves and fine scratches (striations) resulting from this scraping process are often seen on freshly exposed bedrock, and they are important indicators of the direction of ice movement. Erosion and sediment deposition by the ice sheet combined to give a streamlined shape to many hills, with their long dimension parallel to the direction of ice flow. Some of these hills (drumlins) are composed of dense glacial sediment (till) plastered under great pressure beneath the ice.

A warming climate forced the ice sheet to start retreating as early as 21,000 calendar years ago, soon after it reached its southernmost position on Long Island (Ridge, 2004). The edge of the glacier withdrew from the continental shelf east of Long Island and reached the present position of the Maine coast by about 16,000 years ago (Borns and others, 2004). Even though the weight of the ice was removed from the land surface, the Earth's crust did not immediately spring back to its normal level. As a result, the sea flooded much of southern Maine as the glacier retreated to the northwest. Ocean waters extended far up the Kennebec and Penobscot valleys, reaching present elevations of up to 420 feet in the central part of the state.

Great quantities of sediment washed out of the melting ice and into the sea, which was in contact with the retreating glacier margin. Sand and gravel accumulated as deltas and submarine fans where streams discharged along the ice front, while the finer silt and clay dispersed across the ocean floor. The shells of clams, mussels, and other invertebrates are found in the glacial-marine clay that blankets low land areas of southern Maine. Ages of these fossils tell us that ocean waters covered parts of Maine until about 13,000 years ago. The land rebounded as the weight of the ice sheet was removed, forcing the sea to retreat.


References Cited

- Borns, H. W., Jr., Doner, L. A., Dorion, C. C., Jacobson, G. L., Jr., Kaplan, M. L., Kretz, K. J., Lowell, T. V., Thompson, W. B., and Weddle, T. K., 2004, The deglaciation of Maine, U.S.A., in Ehlers, J., and Gibbard, P. L., eds., *Quaternary Glaciations - Extent and Chronology, Part II: North America*, Amsterdam, Elsevier, p. 89-109.
- Davis, R. B., and Jacobson, G. L., Jr., 1985, Late-glacial and early Holocene landscapes in northern New England and adjacent areas of Canada: *Quaternary Research*, v. 23, p. 341-368.
- Ridge, J. C., 2004, The Quaternary glaciation of western New England with correlations to surrounding areas, in Ehlers, J., and Gibbard, P. L., eds., *Quaternary Glaciations - Extent and Chronology, Part II: North America*, Amsterdam, Elsevier, p. 169-199.
- Stone, B. D., and Borns, H. W., Jr., 1986, Pleistocene glacial and interglacial stratigraphy of New England, Long Island, and adjacent Georges Bank and Gulf of Maine, in Sibera, V., Bowen, D. Q., and Richmond, G. M. (editors), *Quaternary glaciations in the northern hemisphere: Quaternary Science Reviews*, v. 5, p. 39-52.



SOURCES OF INFORMATION

Surficial geologic mapping of the Camden quadrangle was conducted by Woodrow B. Thompson during the 2009 field season. Funding for this work was provided by the U.S. Geological Survey STATEMAP program and the Maine Geological Survey, Department of Conservation.



Topographic base from U.S. Geological Survey Camden quadrangle, scale 1:24,000 using standard U.S. Geological Survey topographic map symbols.

The use of industry, firm, or local government names on this map is for location purposes only and does not implicate responsibility for any present or potential effects on the natural resources.

- Ha** Stream alluvium - Sand, gravel, and silt deposited on flood plains of modern streams. May include some wetland deposits.
- Hms** Marine shoreline deposits - Sand and gravel on modern ocean beaches.
- Hw** Wetland deposits - Peat, muck, silt, and clay in poorly drained areas.
- Pmn** Marine nearshore deposits - Sandy to gravelly sediments formed in late-glacial time when marine processes reworked glacial deposits during regression of the sea.
- Pp** Presumpscot Formation - Glaciomarine silt, clay, and sand deposited on the late-glacial sea floor. This map unit overlies the irregular surface of glacial till in a complex manner, so it is likely to include areas of till exposed at the ground surface.
- Pmd** Glaciomarine delta - Sand and gravel deposited into the sea and built up to the ocean surface. Formed at the glacier margin during recession of the most recent (late Wisconsinan) ice sheet. Elevation of boundary between topset and foreset beds (T/F contact) in each delta indicates the position of sea level when the delta was deposited.
- Pmdwr** West Rockport delta. T/F contact not exposed.
- Pt** Till - Loose to very compact, poorly sorted, massive to weakly massive mixture of sand, silt, and gravel-size rock debris deposited by glacial ice. Boulders are commonly scattered across the ground surface. This map unit locally includes lenses of water-laid sand and gravel, as well as patches of overlying Presumpscot Formation (Pp).
- Ptd** Thin drift, undifferentiated - Area of thin patchy sediment cover on bedrock, in which exposure is generally poor and the sediments may include till, Presumpscot Formation, and/or marine nearshore deposits. This unit occurs mainly in coastal areas and islands where bedrock outcrops are extensive.
- af** Artificial fill - Variable mixtures of earth, rock, and/or man-made materials used as fill for roads, and for docks, buildings, and parking lots on waterfront in Rockport Harbor.
- 185** Glacial striation locality - Arrow shows ice-flow direction(s) inferred from striations on bedrock. Dot marks point of observation. Number is azimuth (in degrees) of flow direction. At sites where two or more sets of striations are present and relative ages could be determined, the flinged arrows indicate the older flow directions.
- Moraine ridge - Line shows inferred crest of moraine ridge deposited along the retreating margin of the most recent glacial ice sheet. These moraines are composed mostly of till, but may also include sand and gravel.
- Marine shoreline - Line indicates a former marine shoreline that is 270 ft higher than present-day sea level. The shoreline is marked by the boundary between the till-covered hillside to the west and a gently sloping terrace underlain by marine nearshore sediments (silt, sand, and gravel). Wave-washed bedrock also occurs in this area. The shoreline marks the upper limit of marine submergence in the Camden area during late-glacial time (Berry and others, 2009; Kelley and others, 2000).
- Meltwater channel - Channel eroded by glacial meltwater stream. Arrow shows inferred direction of water flow.
- ↗** Glacially streamlined hill - Symbol shows trend of long axis, which is parallel to former glacial ice-flow direction.

- USES OF SURFICIAL GEOLOGY MAPS**
- A surficial geology map shows all the loose materials such as till (commonly called hardpan), sand and gravel, or clay, which overlie solid bedrock (bedrock). Bedrock outcrops and areas of abundant bedrock outcrops are shown on the map, but varieties of the bedrock are not distinguished (refer to bedrock geology map). Most of the surficial materials are deposits formed by glacial and deglacial processes during the last stage of continental glaciation, which began about 25,000 years ago. The remainder of the surficial deposits are the products of postglacial geologic processes, such as river floodplains, or are attributed to human activity, such as fill or other land-modifying features.
- The map shows the areal distribution of the different types of glacial features, deposits, and landforms as described in the map explanation. Features such as striations and moraines can be used to reconstruct the movement and position of the glacier and its margin, especially as the ice sheet melted. Other ancient features include shorelines and deposits of glacial lakes or the glacial sea, now long gone from the state. This glacial geologic history of the quadrangle is useful to the larger understanding of past earth climate, and how our region of the world underwent recent geologically significant climatic and environmental changes. We may then be able to use this knowledge in anticipation of future similar changes for long-term planning efforts, such as coastal development or waste disposal.
- Surficial geology maps are often best used in conjunction with related maps such as surficial materials maps or significant sand and gravel aquifer maps for any one wanting to know what lies beneath the land surface. For example, these maps may aid in the search for water supplies, or economically important deposits such as sand and gravel for aggregate or clay for bricks or pottery. Environmental issues such as the location of a suitable landfill site or the possible spread of contaminants are directly related to surficial geology. Construction projects such as locating new roads, excavating foundations, or siting new homes may be better planned with a good knowledge of the surficial geology of the site. Refer to the list of related publications below.

OTHER SOURCES OF INFORMATION

- Thompson, W. B., 2010, Surficial materials of the Camden quadrangle, Maine: Maine Geological Survey, Open-File Map 10-5.
- Thompson, W. B., 1979, Surficial geology handbook for coastal Maine: Maine Geological Survey, 68p. (out of print).
- Thompson, W. B., and Borns, H. W., Jr., 1985, Surficial geologic map of Maine: Maine Geological Survey, scale 1:500,000.
- Berry, H. N., IV, Thompson, W. B., Rensch, D. N., and Weddle, T. K., 2009, Geology of the Camden Hills Area [Maine]: Geological Society of Maine, guidebook for summer field trip, 26 p.
- Kelley, J. T., Thompson, W. B., Dab, J., and Bellup, D. F., 2000, Quaternary stratigraphy, sea-level change and bluff erosion, West Penobscot Bay, Maine, in Yates, M. G., Lux, D. R., and Kelley, J. T. (editors), *Guidebook to field trips in coastal and east-central Maine: New England Intercollegiate Geological Conference, 92nd annual meeting, October 6-8, 2000, Orono, Maine*, p. 238-255.



Figure 1. Mt. Battie (left) and Mt. Megunticook overlook Camden in the foreground. These glacially sculpted hills are composed of hard quartz-rich bedrock of probable Cambrian age.



Figure 2. Outcrop of quartzite conglomerate on Mt. Battie. This distinctive rock unit commonly shows polished surfaces and striations caused by glacial abrasion.



Figure 3. Outcrop of Coombs Limestone next to Beachamp Point Road on east side of Rockport Harbor. Postglacial weathering of the ledge has emphasized the prominent fold structures by etching out the softer rock layers.



Figure 4. Erosion of the sediment cover due to rising sea level has exposed the glacial bedrock surface in many places along the Maine coast. This example on the east side of Rockport Harbor shows glacial grooves in the foreground and a glacially sculpted "walldack" form just beyond the knapsack. The grooves indicate ice flow toward the south-southeast (172°, i.e. from right to left as seen in photo).



Figure 5. View looking west at sloping surface of glacial bedrock outcrop, west of U.S. Route 1 in Rockport. The striations record two directions of ice flow. The red pencil is parallel to the younger and dominant set of striations trending 141° (SE). The blue pen parallels older striations trending 160° (SSE). The latter set has been preserved in a few broad grooves that were not completely removed by the younger phase of ice flow.



Figure 6. A stony heterogeneous sediment called "till" was released from melting glacial ice over much of the Camden Quadrangle. Boulders scattered across the ground surface often indicate the presence of till, as seen in this road cut.



Figure 7. Many low areas around Camden are underlain by clay, silt, and sand deposited on the sea floor during the period of marine submergence that immediately followed glacial retreat. This muddy sediment is called the Presumpscot Formation, known locally as "blue clay" where fresh and unweathered. Natural exposures of the clay are not readily seen in the map area (the one shown here is actually in nearby Waldoboro) and are most likely to be found along stream banks and the ocean shore. The embedded pebble in the upper part of the photo probably dropped to the ocean bottom from a floating iceberg.



Figure 8. At least the gravely upper part of this well-stratified deposit is thought to have resulted from marine erosion and reworking of glacial till on a nearby hillside in Camden. It is mapped as unit Pmn (marine nearshore deposit). The deeper and sandier portion may have washed into the sea directly from melting glacial ice.