



Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF MAINE.

Bravo® ZN

EPA Reg. No. 50534-204

EPA SLN No. ME-100001

For Control of Late Blight (*Phytophthora infestans*) for Long-Season Potatoes

KEEP OUT OF REACH OF CHILDREN WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label. This label must be in the possession of the user at the time of pesticide application.

CROP	DISEASES (Pathogen)	Pts. Product./A (lbs. a.i./A)	APPLICATION DIRECTIONS
Potato (Long-season varieties)	Late Blight (<i>Phytophthora infestans</i>)	1½ to 2 1/8 (0.75 to 1.125)	<p>Begin applying at 5 to 10 day intervals when Late Blight forecasting measures 18 disease severity values (DSV).</p> <p>Increase water spray volume as canopy density increases. Use the highest rate and shortest interval when plants are rapidly growing and disease conditions are severe.</p> <p>Bravo Zn may be applied through sprinkler irrigation equipment (solid set, portable wheel move, motorized lateral move or center pivot systems only).</p> <p>Do not exceed a 10 day interval between applications when using chemigation. See <i>Application and Calibration Techniques for Chemigation</i> on Bravo Zn label.</p>

Specific Use Restrictions: Do not apply more than 30½ pts. Bravo Zn (16 lbs. a.i.) per acre on long-season varieties of potato during each growing season. Do not apply within 7 days of harvest.

Bravo® is a trademark of a Syngenta Group Company

Section 24(c) Registrant:
GB Biosciences
Greensboro, NC

ME0204015AA0510