



Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF MAINE

Callisto® Herbicide

**EPA REG. NO. 100-1131
EPA SLN NO. ME-120001**

This label expires and must not be distributed or used in accordance with this SLN registration after June 1, 2017

FOR WEED CONTROL IN LOWBUSH BLUEBERRY IN CROP-BEARING YEARS

<i>Active Ingredient:</i>	
Mesotrione: (CAS No. 104206-82-8)	40.0%
<i>Other Ingredients:</i>	60.0%
<i>Total:</i>	100.0%

Callisto contains 4 lbs of active ingredient mesotrione per gallon.

CAUTION

KEEP OUT OF REACH OF CHILDREN

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR PEST CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

General Information

Callisto is used on lowbush blueberry for control or suppression of many broadleaf weeds.

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, and precautions including statements pertaining to the Worker Protection Standard, on the EPA-registered Callisto label.

This labeling must be in the possession of the user at the time of herbicide application.

Specific Use Directions- Lowbush Blueberry

Apply Callisto as a broadcast spray at a rate of 4.0 fl. oz/A to lowbush blueberry for control or suppression of common lambsquarters, redroot pigweed, velvetleaf, wild mustard, spreading dogbane, blue violet, sheep sorrel, goldenrod and common ragweed. The application of Callisto can be made prior to weed emergence or after weed emergence but before weeds reach 5" in height.

The use of a non-ionic surfactant (NIS) type adjuvant at 0.25% v/v (1 qt/100 gallons of spray volume) is recommended.

Applications of Callisto during dry weather conditions and/or temperatures above 85 degrees can cause injury to lowbush blueberries. Applications of Callisto can cause yellowing or necrosis of leaves and under severe conditions, leaf drop may occur especially on "Sourtop" variety blueberries.

Restrictions:

1. Make only one application per year.
2. The application of Callisto must be made prior to lowbush blueberry bloom.
3. Do not harvest within 60 days of application.
4. Do not apply by air.

Callisto® trademark of a Syngenta Group Company

24(c) registrant:

Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, NC 27419-8300

ME1131018BA0512