Jeanne M. Lambrew, Ph.D. Commissioner




Maine Monthly Pertussis Surveillance Report

July 2019

Date of Report: August 1, 2019

This update summarizes cases of pertussis reported to the Maine Center for Disease Control from January 1st through July 31st, 2019. There were 22 pertussis cases reported statewide in June, compared to 31 reported cases in 2018 and a 5-year median of 24 during that same time (see below graph). Six counties had rates of pertussis above the state rate: Cumberland, Hancock, Knox, Lincoln, Waldo and Washington. Year to date, most cases were in kids between the ages of 11-19 (52.6%), the average age was 15.2 years and the median age was 14 years.

There were no outbreaks reported in July.



- The clinical case definition and surveillance case definitions of pertussis are different, so the reported number of cases may seem lower than expected
- All data presented is preliminary and subject to change

Jeanne M. Lambrew, Ph.D. Commissioner



Table 1. 2019 Monthly Case Count, YTD Case Count and YTDIncidence Rate per 100,000

County	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Total Cases (YTD)	Incidence Per 100,000 (YTD)
Aroostook	0	0	0	0	0	0	0						0	0.0
Androscoggin	4	1	3	0	0	2	0						10	9.3
Cumberland	3	4	17	11	13	18	7						73	25.2
Franklin	0	0	0	0	0	0	0						0	0.0
Hancock	16	9	5	4	8	4	6						52	95.1
Kennebec	0	0	2	1	2	0	0						5	4.2
Knox	2	4	1	1	8	8	3						27	67.7
Lincoln	2	4	2	1	0	1	1						11	32.4
Oxford	0	1	0	1	1	0	1						4	7.0
Penobscot	1	0	2	0	0	0	0						3	2.0
Piscataquis	0	0	0	0	0	0	0						0	0.0
Sagadahoc	1	0	0	0	0	0	0						1	2.8
Somerset	0	1	0	0	0	0	0						1	2.0
Waldo	10	11	4	5	3	1	1						35	89.4
Washington	3	2	2	3	0	0	0						10	31.6
York	13	3	3	6	4	2	3						34	16.9
Maine	55	40	41	33	39	36	22						266	20.0

Table 2. Pertussis cases by age group- Maine, Year to Date 2019

Age Group	# of Cases	Percent of Cases			
< 6 Months	4	1.5%			
6-11 Months	10	3.8%			
1-6 Years	48	18.0%			
7-10 Years	28	10.5%			
11-19 Years	140	52.6%			
≥ 20 Years	36	13.5%			

Demographics: Average age: 15.2years, Median age: 14 years. Min 1 month, Max 71 years

50% Female, 50% Male

- The clinical case definition and surveillance case definitions of pertussis are different, so the reported number of cases may seem lower than expected
- All data presented is preliminary and subject to change

Janet T. Mills Governor

Jeanne M. Lambrew, Ph.D. Commissioner



Maine Department of Health and Human Service Maine Center for Disease Control and Preventio 11 State House Statio 286 Water Stree Augusta, Maine 04333-001 Tel; (207) 287-8016; Fax (207) 287-905 TTY: Dial 711 (Maine Relay

Table 3. Pertussis Outbreaks by County and Facility
Type- Maine, Year to Date 2019CountyFacility TypeSchool (K-12)DaycareOtherCumberland41Hancock11Lincoln22

Pertussis Prevention:

2

York

- The best way to prevent pertussis (whooping cough) among babies, children, teens, and adults is to get vaccinated.
 - For more information on pertussis vaccines visit <u>https://www.cdc.gov/pertussis/vaccines.html</u>
- Pertussis spreads by coughing and sneezing while in close contact with others, who then breathe in the bacteria. Practice good hand hygiene to prevent the spread of respiratory illnesses. To practice good hygiene, you should:
 - Cover your mouth and nose with a tissue when you cough or sneeze.
 - Put your used tissue in the waste basket.
 - Cough or sneeze into your upper sleeve or elbow, not your hands, if you don't have a tissue.
 - Wash your hands often with soap and water for at least 20 seconds or use an alcohol-based hand rub if soap and water are not available.

- The clinical case definition and surveillance case definitions of pertussis are different, so the reported number of cases may seem lower than expected
- All data presented is preliminary and subject to change