

2008 MAINE STATE PROFILE of SELECTED PUBLIC HEALTH INDICATORS

Maine Center for Disease Control and Prevention/DHHS

DEMOGRAPHICS	DISTRICT									COMPARISONS	
	Aroos. ± Margin of Error	Central ± Margin of Error	Cumb. ± Margin of Error	DnEast ± Margin of Error	MdCoast ± Margin of Error	Penquis ± Margin of Error	Wash. ± Margin of Error	West ± Margin of Error	York ± Margin of Error	MAINE State ± Margin of Error	UNITED STATES
US Census: Official Population Estimates											
Total Population [2006]	73,008	173,317	274,598	87,085	151,882	164,765	33,288	194,687	202,232	1,321,574	299,398,484
Total Percent of Maine Population [2006]	5.5	13.1	20.8	6.6	11.5	12.5	2.5	14.7	15.3	100%	n/a
Population Density (people per square mile) [2006]	10.9	36.2	328.7	21.0	84.1	22.4	13.2	45.9	204.1	42.8	79.6 [2000]
Median Age (*see Tech notes) [2006]	43.0	41.1*	40.7	42.9*	42.3*	41.9*	42.7	40.0*	41.1	41.1	36.4
Selected Ages (US Census Official Population Estimates) [2006]											
<5 Years of Age (percent and count)	4.8 (3,478)	5.2 (9,059)	5.6 (15,254)	5.0 (4,379)	5.1 (7,731)	5.3 (8,701)	5.0 (1,767)	5.5 (10,745)	5.4 (10,898)	5.3 (70,245)	6.8 (20,417,636)
65 Years & Older (percent and count)	17.4 (12,673)	14.5 (25,175)	13.6 (37,237)	16.6 (14,462)	15.7 (23,912)	14.0 (23,007)	17.0 (5,902)	14.4 (28,007)	13.9 (28,166)	14.1 (192,639)	12.4 (37,260,352)
Race/Ethnicity Counts (one race alone or in combination with one or more other races, except where noted.** US Census Official Population Estimates) *see Tech notes [2006]											
White (one race alone or in combination)	71,059	170,767	264,065	84,587	149,851	160,820	31,591	190,829	198,734	1,290,712 (97%)	81.4%
Black (one race alone or in combination)	481	1,209	5,772	553	1,111	1,433	237	2,624	1,681	14,864 (1.1%)	13.4%
American Indian & Alaskan Native (one race alone or in combination)	1,419	1,601	2,115	2,194	1,322	2,118	1,629	1,609	1,175	13,553 (1%)	1.5%
Asian (one race alone or in combination)	633	1,438	5,413	736	959	1,725	232	1,747	2,149	14,800 (1.1%)	5.0%
Native Hawaiian & Other P.I. (one race alone or in combination)	19	97	310	29	51	126	7	94	80	806 (<1%)	0.3%
Hispanic (of any race)**	636	1,400	3,891	827	1,424	1,369	392	1,965	2,017	13,529 (1%)	14.8%
Non-Hispanic (total)**	72,372	171,917	270,707	86,258	150,458	163,396	32,896	192,722	200,215	1,308,045 (99%)	85.1%
Franco-American [2000]	25,374	46,580	47,185	9,083	19,132	46,580	3,383 [2006]	58,457	52,868	305,259 (22%)	0.8%

2008 MAINE STATE PROFILE of SELECTED PUBLIC HEALTH INDICATORS

Maine Center for Disease Control and Prevention/DHHS

MORE DEMOGRAPHICS: Percent and Count	DISTRICT									COMPARISONS	
	Aroos. ± Margin of Error	Central ± Margin of Error	Cumb. ± Margin of Error	Dneast ± Margin of Error	Mdcoast ± Margin of Error	Penquis ± Margin of Error	Wash. ± Margin of Error	West. ± Margin of Error	York ± Margin of Error	MAINE State ± Margin of Error	UNITED STATES
Median Annual Household Income [2004]	\$32,629	\$36,147	\$49,870	\$33,834	\$41,690	\$34,717	\$29,087	\$ 36,670	\$ 48,363	\$ 41,287	\$ 44,334
Families Living in Poverty (all ages, percent) [2004]	14.9 (±0.3) 10,918	13.1 (±0.2) 22,497	9.0 (±0.1) 24,708	12.6 (±0.2) 10,943	10.7 (±0.2) 16,142	13.2 (±0.2) 21,715	17.4 (±8.2) 5,701	12.4 (±0.2) 23,915	9.0 (±0.1) 17,963	11.3 (±0.1) 148,801	11.5
Children on Free or Reduced Lunch Program (percent of enrolled school children) [2005]	47.7 (±1.2) 5,551	41.0 (±0.8) 11,613	25.9 (±0.5) 10,483	41.9 (±1.1) 5,037	35.9 (±0.8) 7,471	39.1 (±0.8) 10,068	55.5 2,772	45.8 (±0.7) 13,466	27.6 (±0.7) 7,547	36.4 (±0.3) 71,236	17.5
Adults with Lifetime Educational Attainment Less Than High School (percent) [2000]	23.1 (±0.4) 11,868	16.1 (±0.3) 18,378	9.8 (±0.1) 17,900	15.3 (±0.3) 9,185	13.0 (±0.2) 13,146	14.9 (±0.2) 16,073	20.1 4,729	18.6 (±0.2) 23,561	13.5 (±0.2) 17,177	14.6 (±0.1) 127,419	9.4 (±0.1)
Single-Parent Households with Children <18 years (percent) [2000]	7.7 (±0.3) 2,323	10.4 (±0.2) 7,007	8.4 (±0.2) 9,117	8.8 (±0.3) 3,157	9.0 (±0.2) 5,372	9.5 (±0.2) 6,197	5.9 839	10.4 (±0.2) 7,887	9.1 (±0.2) 6,788	9.2 (±0.1) 47,848	7.2
Householders ≥ 65 Living Alone (percent) [2000]	13.1 (±0.4) 3,977	10.5 (±0.2) 7,145	10.2 (±0.2) 11,015	12.2 (±0.3) 4,386	11.0 (±0.4) 6,549	10.4 (±0.2) 6,829	13.1 1,847	10.9 (±0.2) 8,317	9.7 (±0.2) 7,233	10.7 (±0.1) 55,451	9.2
People Who Speak a Language Other Than English (percent of those >5 years old) [2000]	24.1 (±0.3) 16,880	6.8 (±0.1) 10,735	5.9 (±0.1) 14,888	4.3 (±0.1) 3,498	3.5 (±0.1) 4,798	4.4 (±0.1) 6,818	5.4 1,731	11.1 (±0.2) 19,771	9.4 (±0.1) 16,578	7.8 (±0.1) 82,512	17.9
People with Disability Status (percent of those >5 years old) [2000]	24.8 16,924	14.0 24,905	17.1 42,102	20.0 17,830	20.0 38,143	18.0 30,727	27.0 8,543	20.0 38,143	18.9 32,913	20.0 237,910	19.3

2008 MAINE STATE PROFILE of SELECTED PUBLIC HEALTH INDICATORS

Maine Center for Disease Control and Prevention/DHHS

MATERNAL/CHILD HEALTH INDICATORS	Aroos. ± Margin of Error	Central ± Margin of Error	Cumb. ± Margin of Error	Dneast ± Margin of Error	Mdcoast ± Margin of Error	Penquis ± Margin of Error	Wash. ± Margin of Error	West. ± Margin of Error	York ± Margin of Error	MAINE State ± Margin of Error	UNITED STATES	Bench- mark State (healthiest)
Infant Mortality (rate per 1,000 live births) [2001-2005]	6.1 (±2.6)	5.2 (±1.5)	5.2 (±1.2)	4.4 (±2.0)	5.3 (±1.6)	6.0 (±1.6)	4.6 (±3.2)	6.0 (±1.5)	5.8 (±1.4)	5.5 (±0.5)	6.8 [2004]	UT 4.5 [2005]
Live Births with Low Birth Weight <2500 grams (percent of live births) [2006]	6.6 (±1.9)	6.5 (±1.1)	6.8 (±0.8)	6.0 (±1.5)	7.0 (±1.3)	7.3 (±1.2)	7.9 (±2.9)	6.6 (±1.0)	7.4 (±1.2)	6.8 (±0.4)	8.2 [2005]	AK+OR+ WA 6.1 [2005]
Infants Born to Women Receiving First Trimester Prenatal Care (percent) [2006]	90.4 (±2.2)	80.9 (±1.8)	89.7 (±1.1)	86.3 (±2.2)	88.7 (±1.5)	85.3 (±1.7)	82.5 (±3.9)	88.9 (±1.3)	88.3 (±1.4)	87.4 (±0.6)	83.9 [2005]	RI + MA 89.3 [2005]
Teen Births Ages 15-17 (rate per 1,000 female population) [2003-2005]	14.0 (±3.4)	13.7 (±2.2)	8.7 (±1.5)	9.2 (±2.5)	11.8 (±2.2)	11.4 (±2.1)	14.3 (±5.0)	15.1 (±2.2)	7.5 (±1.5)	11.2 (±0.7)	21.4 [2005]	NH 7.0 [2005]
HEALTH & WELLNESS INDICATORS												
Adolescent Smoking Prevalence (percent of 6-12 graders) [2006]	15.3 (±0.6)	14.4 (±0.5)	11.9 (±0.3)	16.0 (±0.6)	14.8 (±0.5)	16.1 (±0.6)	15.5	13.3 (±0.4)	12.5 (±0.4)	13.8 (±0.2)	23.0 [2005]	UT: 7.4 ID: 15.8 [2005] HS only
Adult Smoking Prevalence (percent who are current smokers) [2006]	28.4 (±7.8)	23.1 (±4.7)	16.3 (±3.7)	24.8 (±5.7)	18.2 (±3.1)	26.7 (±4.9)	28.6 (±7.9)	17.6 (±3.9)	21.7 (±5.5)	21.0 (±1.6)	20.1	UT: 9.8 CA: 14.9
High School Youth Overweight or Obese (percent) [2005]	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	25.0	26.3	CO 20.1
Adults Overweight or Obese (percent) [2006]	55.0 (±8.2)	63.9 (±5.7)	52.8 (±5.1)	60.9 (±6.1)	60.9 (±4.1)	64.7 (±5.3)	68.0 (±7.9)	59.4 (±5.3)	64.9 (±5.5)	60.2 (±2.0)	61.6	CO + UT 54.9
Adults Reporting Fair or Poor Health Status in last 30 days (percent) [2006]	15.4 (±5.4)	16.5 (±4.0)	11.0 (±2.9)	16.7 (±4.3)	11.7 (±2.5)	18.7 (±4.0)	21.7 (±6.9)	11.7 (±2.8)	11.0 (±3.5)	13.6 (±1.3)	14.7	MN 10.8

2008 MAINE STATE PROFILE of SELECTED PUBLIC HEALTH INDICATORS

Maine Center for Disease Control and Prevention/DHHS

CHRONIC DISEASE INDICATORS	Aroos. ± Margin of Error	Central ± Margin of Error	Cumb. ± Margin of Error	Dneast ± Margin of Error	Mdcoast ± Margin of Error	Penquis ± Margin of Error	Wash. ± Margin of Error	West. ± Margin of Error	York ± Margin of Error	MAINE State ± Margin of Error	UNITED STATES	Benchmark State (healthiest)
Overall Cancer Incidence (age-adjusted rate per 100,000) [2000-2004]	514.2 (±20.9)	524.5 (±14.6)	500.4 (±11.6)	572.9 (±20.4)	513.4 (±15.0)	553.1 (±15.6)	608.9 [2003-2005]	505.7 (±13.4)	495.3 (±13.4)	517.7 (±5.2)	458.2 [2004]	AZ: 383.3 NM: 409.0 [2004]
Overall Cancer Mortality (age-adjusted rate per 100,000) [2000-2004]	199.0 (±12.7)	206.4 (±9.1)	208.5 (±7.4)	223.3 (±12.6)	200.1 (±9.3)	217.5 (±9.8)	241.6	208.0 (±8.6)	201.5 (±8.6)	207.6 (±3.3)	185.7 [2004]	UT: 139.1 HI: 147.5 [2004]
Lung Cancer Incidence (age-adjusted rate per 100,000) [2000-2004]	84.5 (±8.4)	79.7 (±5.7)	77.3 (±4.6)	88.8 (±8.0)	74.2 (±5.7)	92.9 (±6.4)	102.1 [2003-2005]	82.5 (±5.4)	72.8 (±5.2)	80.6 (±2.1)	67.4 [2004]	UT: 28.3 NM: 45.3 [2004]
Lung Cancer Mortality (age-adjusted rate per 100,000) [2000-2004]	62.4 (±7.2)	58.5 (±4.9)	59.8 (±4.1)	61.7 (±6.7)	57.2 (±5.0)	69.4 (±5.5)	77.2	65.4 (±4.8)	59.0 (±4.7)	61.5 (±1.8)	61.0 [2004]	UT: 26.2 NM: 35.8 [2004]
Colorectal Cancer Incidence (age-adjusted rate per 100,000) [2000-2004]	65.9 (±7.5)	58.7 (±4.9)	54.0 (±3.8)	61.8 (±6.7)	57.7 (±5.0)	67.1 (±5.5)	77.7 [2003-2005]	50.2 (±4.3)	60.7 (±4.8)	58.4 (±1.8)	49.5 [2004]	UT: 37.4 AZ: 39.4 [2004]
Colorectal Cancer Mortality (age-adjusted rate per 100,000) [2000-2004]	24.5 (±4.6)	19.9 (±2.9)	22.2 (±2.5)	23.8 (±4.2)	18.9 (±2.9)	20.5 (±3.1)	27.3	17.2 (±2.5)	20.4 (±2.8)	20.5 (±1.1)	17.9 [2004]	UT: 12.2 HI: 13.4 [2004]
Sigmoidoscopy or Colonoscopy Screening Ever Had by Adults Age 50 and Older (percent) [2006]	53.2 (±10.0)	71.8 (±6.3)	74.3 (±5.1)	54.1 (±7.6)	61.8 (±5.1)	71.8 (±6.3)	38.6 [2000-2006]	60.0 (±6.5)	64.4 (±6.9)	64.6 (±2.4)	57.1 (median % of states responding)	RI 69.2
Female Breast Cancer Incidence (age-adjusted rate per 100,000) [2000-2004]	116.7 (±14.2)	136.0 (±10.3)	134.8 (±8.2)	136.0 (±10.3)	134.6 (±10.7)	136.7 (±10.7)	127.0 [2003-2005]	127.5 (±9.4)	132.1 (±9.5)	132.5 (±3.6)	117.7 [2004]	AZ:102.9 ID: 105.1 [2004]
Female Breast Cancer Mortality (age-adjusted rate per 100,000) [2000-2004]	16.9 (±5.3)	23.0 (±4.2)	25.1 (±3.5)	23.0 (±4.2)	17.3 (±3.8)	26.5 (±4.7)	31.7	25.7 (±4.2)	23.6 (±4.0)	23.7 (±1.5)	24.4 [2004]	HI: 15.6 AK: 18.7 [2004]

2008 MAINE STATE PROFILE of SELECTED PUBLIC HEALTH INDICATORS

Maine Center for Disease Control and Prevention/DHHS

CHRONIC DISEASE INDICATORS (cont'd)	Aroos. ± Margin of Error	Central ± Margin of Error	Cumb. ± Margin of Error	Dneast ± Margin Of Error	Mdcoast ± Margin of Error	Penquis ± Margin of Error	Wash. ± Margin of Error	West. ± Margin of Error	York ± Margin of Error	MAINE State ± Margin of Error	UNITED STATES	Bench- mark State (healthiest)
Mammogram in Past Two Years Among Women 40 and Older (percent) [2006]	85.2 (±7.3)	84.7 (±5.5)	85.1 (±4.3)	75.5 (±8.0)	77.3 (±5.1)	84.7 (±5.5)	75.5 (±10.7)	83.3 (±5.5)	81.5 (±5.7)	82.0 (±2.0)	76.5 (median % of states responding)	MA 84.8
Pap Smear in Past 3 Years Among Women 18 and Older (percent) [2006]	90.9 (±5.7)	89.6 (±5.3)	91.6 (±3.3)	88.1 (±5.5)	83.7 (±4.5)	89.6 (±5.3)	90.0 (±6.3)	92.6 (±4.3)	90.4 (±4.3)	89.1 (±1.6)	84.0 (median % of states responding)	ME 89.1
Prostate Cancer Incidence (age-adjusted rate per 100,000) [2000-2004]	174.7 (±18.3)	182.0 (±13.2)	161.1 (±10.2)	190.8 (±17.6)	173.6 (±13.0)	173.8 (±13.4)	179.1 [2003-2005]	192.8 (±12.6)	159.2 (±11.6)	174.5 (±4.6)	145.3 [2004]	AZ: 109.7 MO: 117.2 [2004]
Prostate Cancer Mortality (age-adjusted rate per 100,000) [2000-2004]	21.7 (±7.0)	28.1 (±5.7)	29.8 (±4.8)	31.3 (±7.7)	33.8 (±6.2)	28.9 (±6.3)	29.4	25.0 (±5.0)	27.2 (± 5.3)	28.5 (± 2.1)	25.4 [2004]	HI: 18.9 FL: 22.0 [2004]
Major CVD Deaths (rate per 100,000) [2005] ICD-10 codes 100-178	286.8 (±33.0)	254.0 (±21.8)	204.6 (±15.5)	262.8 (±29.0)	236.8 (±21.2)	282.2 (±24.1)	454.4 [2001-2005]	251.4 (±20.3)	215.5 (±18.8)	242.0 (±7.6)	286.6 [2004]	MN 210.1 [2004]
Stroke Hospitalizations (rate per 10,000) [2005]	24.9 (±3.1)	18.8 (±1.9)	21.7 (±1.6)	19.5 (±2.5)	19.5 (±2.0)	20.6 (±2.0)	19.4	22.2 (±1.9)	17.8 (±1.6)	20.7 (±0.7)	n/a	n/a
Acute Myocardial Infarction Hospitalizations (rate per 10,000) [2005]	57.1 (±4.7)	37.7 (±2.7)	19.6 (±1.5)	41.1 (±3.7)	24.2 (±2.0)	31.9 (±2.5)	38.7	22.9 (±2.0)	22.0 (±1.9)	29.2 (±0.8)	n/a	n/a
High Blood Pressure Among Adults (percent) [2005]	28.7 (±6.9)	25.3 (±4.5)	18.7 (±3.1)	28.9 (±5.5)	25.5 (±3.5)	29.6 (±4.7)	32.3 (±7.7)	25.0 (±3.9)	28.5 (±4.7)	25.4 (±1.6)	25.5 (median % of states responding)	UT 18.4
High Cholesterol Among Adults (percent) [2005]	35.0 (±7.8)	36.4 (±5.1)	29.3 (±4.1)	39.8 (±6.3)	36.5 (±4.3)	39.1 (±5.5)	44.4 (±9.2)	42.9 (±5.1)	36.4 (±5.5)	36.4 (±2.0)	35.6 (median % of states responding)	LA 30.3

2008 MAINE STATE PROFILE of SELECTED PUBLIC HEALTH INDICATORS

Maine Center for Disease Control and Prevention/DHHS

CHRONIC DISEASE INDICATORS (cont'd)	Aroos. ± Margin of Error	Central ± Margin of Error	Cumb. ± Margin of Error	Dneast ± Margin of Error	Mdcoast ± Margin of Error	Penquis ± Margin of Error	Wash. ± Margin of Error	West. ± Margin of Error	York ± Margin of Error	MAINE State ± Margin of Error	UNITED STATES	Benchmark State (healthiest)
Diabetes Prevalence Among Adults (non-gestational; percent) [2004-2006]	11.4 (±2.7)	7.7 (±1.6)	6.0 (±1.2)	7.1 (±1.6)	6.7 (±1.2)	8.7 (±1.6)	8.4 (±2.7)	6.4 (±1.4)	7.1 (±1.6)	7.3 (±0.6)	7.5 [2006]	CO 5.3 [2006]
Diabetes Hospitalizations (age-adjusted per 10,000) [2005]	13.5 (±2.4)	11.3 (±1.5)	9.6 (±1.1)	8.5 (±1.8)	11.3 (±1.6)	12.3 (±1.6)	8.2	11.5 (±1.5)	7.5 (±1.2)	10.5 (±0.5)	n/a	n/a
Adults with Diabetes Who Have Received a Hemoglobin A1c Test at Least Once Yearly (percent) [2004-2006]	90.2 (±9.8)	94.8 (±5.1)	93.1 (±4.3)	93.2 (±5.5)	88.9 (±5.3)	88.0 (±6.5)	92.7 (±7.3) [2002-06]	91.2 (±5.1)	95.6 (±3.3)	91.9 (±2.0)	n/a	MO 95.5 [2006]
Adults With Diabetes Who Have Taken a Diabetes Management Course (percent) [2004-2006]	56.1 (±11.4)	63.2 (±7.8)	60.3 (±8.2)	57.0 (±11.8)	60.4 (±8.2)	59.6 (±8.4)	n/a	54.9 (±8.2)	50.7 (±9.2)	58.1 (±3.1)	n/a	MN 77.4 [2006]
Adults with Asthma (percent) [2006]	10.4 (±4.1)	9.1 (±3.1)	8.2 (±2.5)	11.8 (±3.7)	10.5 (±2.5)	12.4 (±3.3)	12.1 (±5.3)	8.7 (±3.1)	8.3 (±2.7)	9.6 (±1.2)	8.5	LA 5.9
Child and Youth Asthma (<18 years old, percent) [2003]	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	10.7 (±1.5)	8.9 [2005]	ID 5.0 [2003]
Asthma Emergency Department Visits (age-adjusted rate per 10,000) [2004]	101.0 (±2.4)	83.7 (±2.6)	51.6 (±1.7)	86.3 (±3.8)	53.3 (±2.3)	71.5 (±2.5)	113.0	77.3 (±1.6)	44.9 (±1.8)	66.1 (±1.4)	64.0	n/a
Adults With a Routine Dental Visit in Past Year (percent) [2006]	61.2 (±7.8)	65.4 (±5.3)	75.4 (±5.1)	69.7 (±5.5)	69.8 (±3.7)	66.9 (±5.1)	64.9 (±7.8)	70.7 (±4.5)	74.4 (±4.9)	70.2 (±1.8)	70.3	CT 80.5

2008 MAINE STATE PROFILE of SELECTED PUBLIC HEALTH INDICATORS

Maine Center for Disease Control and Prevention/DHHS

ENVIRONMENTAL HEALTH INDICATORS	DISTRICT									MAINE State ± Margin of Error	UNITED STATES
	Aroos. ± Margin of Error	Central ± Margin of Error	Cumb. ± Margin of Error	Dneast ± Margin Of Error	Mdcoast ± Margin of Error	Penquis ± Margin of Error	Wash. ± Margin of Error	West. ± Margin of Error	York ± Margin of Error		
Emergency Department Visits for Carbon Monoxide Poisoning (rate per 100,000) [2001-2005]	15.9 (±2.3)	6.9 (±1.9)	4.7 (±1.2)	5.2 (±4.9)	6.3 (±1.9)	6.9 (±1.9)	4.5 (±1.6)	8.0 (±1.9)	6.5 (±1.7)	6.9 (±0.7)	n/a
Elevated Blood Lead Levels Among Screened 1-Year Old Children (percent) [2005-2006]	0.3 (±0.4)	1.9 (±0.8)	1.1 (±0.4)	1.0 (±0.7)	1.9 (±0.8)	1.6 (±0.6)	n/a	1.7 (±0.6)	1.4 (±0.5)	1.4 (±0.2)	1.58 (<72 mos. Old) [2005]
Housing Units in Structures Built <1950 (numbers, representing high risk for lead) [2000]	15,244	29,569	45,159	20,370	29,527	27,996	8,338 (38.0%)	37,155	28,112	233,187 (35.8%)	22.3%
Homes with Private Wells Tested for Arsenic (percent) [2003]	n/a	45.6 (±9.7)	n/a	n/a	36.6 (±9.4)	n/a	51.5 (±5.3)	47.1 (±9.4)	n/a	44.6 (±3.9)	n/a
District Community Water Systems Meeting all Health Based Standards (percent) [2006]	78	80	87	76	80	94	86	84	65	80	n/a
District Community Water Systems with Source Water Protection in Place (percent) [2006]	74	78	84	93	88	87	n/a	81	84	84	n/a

2008 MAINE STATE PROFILE of SELECTED PUBLIC HEALTH INDICATORS

Maine Center for Disease Control and Prevention/DHHS

INFECTIOUS DISEASE INDICATORS	DISTRICT									MAINE State ± Margin of Error	UNITED STATES	Bench- mark State (healthiest)
	Aroos. ± Margin of Error	Central ± Margin of Error	Cumb. ± Margin of Error	Dneast ± Margin of Error	Mdcoast ± Margin of Error	Penquis ± Margin of Error	Wash. ± Margin of Error	West. ± Margin of Error	York ± Margin of Error			
Kindergarteners Exempted from Childhood Vaccination for Philosophical Reasons (count) [2007]	3	43	105	58	60	25	2	36	43	373	n/a	n/a
Children Immunized with the 4:3:1:3:3:1 Vaccination Series by 24 Months of Age (percent) [2003-2006] <small>(4:3:1:3:3:1 means at least 4 DTP/DT/DTaP, 3 Polio, 1 Measles-containing, 3 Hib, 3 Hepatitis B, and 1 Varicella vaccine)</small>	90.3 (combined with Downeast)	84.8	77.8	90.3 (combined with Aroostook)	91.0	81.4	n/a	93.6	79.8	87.4	82.0	MA 86.9 [2006]
Influenza Vaccine Past Year for Adults over 65 years (percent) [2006]	69.4 (±13.5)	70.3 (± 9.8)	79.9 (± 6.7)	69.4 (± 13.5)	70.6 (± 6.9)	77.7 (± 8.2)	66.6 (±7.8) [2004-06]	65.6 (±9.2)	69.9 (±9.6)	72.3 (±3.1)	69.6	CO 75.9
Pneumoccal Vaccine Ever Among Adults 65 Years of Age or Older (percent) [2006]	63.4 (±14.7)	73.9 (±10.4)	68.3 (±8.6)	63.4 (±14.7)	66.7 (±7.3)	77.2 (±9.2)	69.3 (±8.9) [2004-06]	63.3 (±10.0)	62.2 (±10.2)	67.9 (±3.5)	66.9	OR 74.7
Chlamydia (total number) [2006]	64	255	652	95	213	303	32	447	280	2,304	n/a	n/a
Late Diagnosis of HIV (number, AIDS diagnosis within 12 months of first HIV diagnosis) [2001-2005]	5	10	31	5	11	9	2	13	12	96	n/a	n/a
Lyme Disease (crude rate per 100,000; 5-year count in parentheses) [2002-2006]	2.4 (9)	7.8 (66)	22.5 (299)	20.8 (89)	24.5 (177)	3.6 (29)	4.4 (7)	6.0 (56)	56.7 (529)	18.8 (1,200)	6.7 [2006]	n/a
Salmonella (crude rate per 100,000; 5-year count in parentheses) [2002-2006]	11.4 (42)	14.2 (119)	13.4 (178)	7.5 (32)	10.5 (76)	8.9 (72)	6.6 (11)	9.7 (91)	11.8 (110)	11.3 (720)	15.5 [2006]	n/a

2008 MAINE STATE PROFILE of SELECTED PUBLIC HEALTH INDICATORS

Maine Center for Disease Control and Prevention/DHHS

INJURY AND VIOLENCE INDICATORS	DISTRICT									MAINE State ± Margin of Error	UNITED STATES	Bench- mark State (healthiest)
	Aroos. ± Margin of Error	Central ± Margin of Error	Cumb. ± Margin of Error	Dneast ± Margin of Error	Mdcoast ± Margin of Error	Penquis ± Margin of Error	Wash. ± Margin of Error	West. ± Margin of Error	York ± Margin of Error			
Motor Vehicle Traffic Crash Deaths (age-adjusted rate per 100,000 and average number per year) [2001-2005]	16.4 (±4.8) 12 avg/yr.	15.0 (±2.6) 26 avg/yr.	9.7 (±1.6) 27 avg/yr.	19.2 (±4.7) 17 avg/yr.	16.7 (±3.0) 25 avg/yr.	13.8 (±2.5) 23 avg/yr.	25.8 (±7.7) 9 avg/yr.	14.9 (±2.4) 30 avg/yr.	12.4 (±2.2) 24 avg/yr.	13.8 (±0.9) 185 avg/yr.	14.5 [2005]	MA 7.8 [2001-05]
Hip Fracture Hospitalizations Among 65+ Year Olds (rate per 100,000 and 5 yr. count) [2001-2005]	707.8 (±65.6) 447	762.7 (±49.0) 932	827.7 (±41.9) 1,497	754.5 (±63.8) 538	739.0 (±49.5) 856	780.6 (±51.6) 878	768.5 (±100.4) 225	745.2 (±45.2) 1,044	649.4 (±43.1) 874	751.3 (±17.5) 7,066	778.4 [2003-05]	n/a
Reported Rapes (rate per 10,000 female population and average number per year) [2001-2005]	2.3 (± 0.5) 16 avg/yr.	5.7 (±0.7) 50 avg/yr.	3.4 (±0.3) 85 avg/yr.	2.3 (±0.6) 10 avg/yr.	3.0 (±0.5) 23 avg/yr.	3.1 (±0.5) 26 avg/yr.	3.4 (±1.2) 6 avg/yr.	7.7 (±0.8) 76 avg/yr.	3.0 (±0.4) 55 avg/yr.	2.8 (±0.1) 340 avg/yr.	n/a	n/a
Domestic Assaults Reported to the Police (rate per 10,000 population and count) [2005]	36.7 (± 4.4) 269	55.8 (± 3.5) 964	40.6 (±2.4) 1,115	22.7 (±3.2) 198	27.0 (±2.6) 411	26.7 (±3.6) 440	23.6 (±5.2) 79	50.7 (±3.2) 986	53.2 (±3.2) 1,076	41.3 (± 1.1) 5,549	n/a	n/a

2008 MAINE STATE PROFILE of SELECTED PUBLIC HEALTH INDICATORS

Maine Center for Disease Control and Prevention/DHHS

SUBSTANCE ABUSE AND MENTAL HEALTH INDICATORS	DISTRICT									MAINE State ± Margin of Error	UNITED STATES	Benchmark State (healthiest)
	Aroos. ± Margin of Error	Central ± Margin of Error	Cumb. ± Margin of Error	Dneast ± Margin of Error	Mdcoast ± Margin of Error	Penquis ± Margin of Error	Wash. ± Margin of Error	West. ± Margin of Error	York ± Margin of Error			
Adults With 14 or More Days of Frequent Mental Distress in the Past Month (percent) [2006]	13.6 (±5.1)	9.7 (±3.0)	11.3 (±3.2)	7.0 (±2.8)	10.3 (±2.6)	12.1 (±3.7)	7.8 (±3.8)	8.7 (±2.6)	8.4 (±3.4)	10.0 (±1.2)	10.2 (±0.6)	MN 6.7
Adults With Current Symptoms of Moderate or Severe Depression (percent) [2006]	5.8 (±3.3)	9.1 (±2.9)	6.5 (±2.7)	7.8 (±3.1)	6.1 (±2.2)	13.3 (±3.9)	9.7 (±4.5)	5.6 (±2.0)	7.0 (±3.1)	7.6 (±1.0)	n/a	ND 5.6 [2006]
Suicide Deaths (age 10 and older, rate per 100,000) [2001-2005]	12.7 (±3.9)	16.2 (±2.9)	12.0 (±2.0)	15.2 (±3.9)	16.9 (±3.1)	15.0 (±2.8)	16.7 (±6.5)	13.9 (±2.9)	13.5 (±2.4)	13.9 (±1.0)	12.6	NY 7.2
Previous 30-Day Alcohol Use Among 9 th -12 th Graders (percent) [2006]	37.3	36.7	41.6	38.0	43.9	40.7	37.2	39.2	42.3	40.3	43.3 [2005]	UT: 15.8 HI: 34.8 [2005]
Adults Who Have Participated in Binge Drinking in the Past 30 Days (percent) [2004]	25.2	27.8	27.3	26.3	27.6	29.8	32.7 [2004-06]	31.4	24.1	27.8	14.4 (median %)	KY 8.6
Binge Drinking Within the Last 2 Weeks Among 9 th -12 th Graders (percent) [2006]	18.4	19.5	22.0	20.8	23.9	22.9	21.2	21.7	22.3	21.6	25.5 Last 30 days [2005]	UT: 8.8 HI: 18.8 [2005]
Previous 30-Day Prescription Drug Misuse Among 9 th -12 th Graders (percent) [2006]	5.4	8.0	7.6	6.5	9.4	8.5	5.2	8.8	9.6	8.2	n/a	n/a
Substance Abuse Admissions (number among all ages) [2006]	1,275	1,204	2,426	1,141	878	1,391	658	901	802	10,018	n/a	n/a

2008 MAINE STATE PROFILE of SELECTED PUBLIC HEALTH INDICATORS

Maine Center for Disease Control and Prevention/DHHS

ACCESS TO CARE INDICATORS	DISTRICT									MAINE State ± Margin of Error	UNITED STATES	Bench- mark State (healthiest)
	Aroos. ± Margin of Error	Central ± Margin of Error	Cumb. ± Margin of Error	Dneast ± Margin of Error	Mdcoast ± Margin of Error	Penquis ± Margin of Error	Wash. ± Margin of Error	West. ± Margin of Error	York ± Margin of Error			
Access to Primary Care Physician (population to physician ratio) [2003]	1,034:1	971:1	759:1	880:1	1,189:1	939:1	828:1 [2000-01]	1,091:1	1,269:1	978:1	1,351:1 [2000]	MA 187.3 per 100,000 [2005]
Adults With No Health Insurance (percent) [2006]	6.3 (±7.4)	12.2 (± 8.5)	5.8 (± 4.4)	15.3 (± 8.5)	11.6 (± 5.8)	12.2 (± 7.8)	14.8 (±5.6)	12.9 (± 6.0)	6.9 (± 6.7)	10.4 (± 2.6)	14.5 (median %)	MN 8.2
PUBLIC HEALTH PREPAREDNESS INDICATORS												
Hospitals with Pandemic Influenza Plan In Process or No Response (percent) [2007]	0	67	14	60	0	25	0	0	0	25	n/a	n/a
Hospitals with Draft Pandemic Influenza Plan Completed (percent) [2007]	0	33	14	40	50	75	100	20	33	35	n/a	n/a
Hospitals with Pandemic Influenza Plan Completed (percent) [2007]	100	0	72	0	50	0	0	80	67	40	n/a	n/a