

RESTRICTED USE PESTICIDE

DUE TO AVIAN AND AQUATIC TOXICITY

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

SUPPLEMENTAL LABELING FOR

INSECT CONTROL ON ORNAMENTALS GROWN OUTDOORS IN NURSERIES: BALSAM FIR/MAINE ONLY

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with all applicable directions, restrictions, and precautions found in the labeling of the federally registered product and its accompanying supplemental labeling.
- Follow all applicable directions, restrictions, and precautions, including statements pertaining to the Worker Protections Standards, on the EPA-registered label.
- This label must be in the possession of the user at the time of application.

DIAZINON AG600 WBC Insecticide

Water-Based Concentrate

EPA Reg. No. 66222-103-34704

ACTIVE INGREDIENT: Diazinon:	% BY WT.
O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate.....	56.0%
INERT INGREDIENTS:	44.0%
	TOTAL 100.0%

ORNAMENTAL STATEMENT

Before treating a large number of ornamental plants with DIAZINON AG600 WBC Insecticide alone or as a tank mixture with any other material, make a test application on a few plants and observe for 7 to 10 days prior to treating large areas to reduce the possibility of plant injury.

INSECT CONTROL ON ORNAMENTALS GROWN OUTDOORS IN NURSERIES

CROP	PEST	RATE	COMMENTS
BALSAM FIR (MAINE ONLY)	Aphids, Bagworms, Carnation Bud Mite, Carnation Shoot Mite, Clover Mite, Cyclamen Mite, Dipterous Leafminer, European Pine Shoot Moth, European Red Mite, Flea Beetles, Holly Bud Moth, Leafhoppers, Obscure Root Weevil, Omnivorous Leaf-tier, Privet Mite, Scale Crawlers (Cottony-cushion, Lecanium, Pine Needle, San Jose, Soft), Thrips, Two-spotted Spider Mite, Juniper Webworm, Whiteflies	½ fl.oz. per 3 gals. of water or 12 ¾ fl.oz. per 100 gals. of water	Apply the recommended rates. Spray when listed pests first appear. Try to spray underside of leaves and penetrate dense foliage.
	Apple-and-Thorn Skeletonizer, Cotoneaster Webworm, European Pine Sawfly, Fall Webworm, Hemlock Chermes, Oak Looper, Obliquebanded Leafroller, Pear Slug, Tent Caterpillar	1 ¼ fl.oz. per 3 gals. of water or 25.6 fl.oz. per 100 gals. of water	
	Mimosa Webworm	2 ½ teaspoons in 3 gals. of water or 13 ¼ fl.oz. in 100 gals. of water	Apply as a thorough foliar spray when adults first appear.

**BALSAM FIR
(MAINE ONLY)
(CONTINUED)**

- Do not use on Ferns, Poinsettia, Hibiscus, Papaya, Pilea, and Gardenia, since injury to the plants may occur.
- The maximum application rate on all ornamentals is 25.6 ounces per acre.
- Make a maximum of one application per crop cycle.
- The REI is 7 days for flowers and other commercial ornamentals grown for cutting.
- The REI is 2 days for all other commercial ornamentals.
- Application is permitted only on commercial ornamentals grown outdoors in nurseries.

Manufactured for:
Loveland Products, Inc.
PO Box 1286
Greeley, Colorado 80632-1286