

TOP FORAGE PLANTS FOR A BEE GARDEN IN MAINE Part I

2013 Edition, Year 2 of 5 Contact: Alison C. Dibble, Ph.D. adibble2@gmail.com

Bees are our most effective pollinators, and flowers are food for bees. To support native bees and honey bees, provide a succession of flowers through the entire growing season. Plants differ in their attraction to bees, and the differences could be subtle. In a University of Maine experiment at four gardens in Old Town, Jonesboro, and Blue Hill, researchers count insects that land on flowers in good weather during three one-minute observations per plant. To date we have tested more than 60 species of native wildflowers, shrubs, bedding plants, cover crops, and herbs. These easy-to-grow bee plants are likely to be successful in your garden:



Anise hyssop, *Agastache foeniculum*, attracts bumble bees. July-Aug. Annual, self sows.



Butterfly milkweed, *Asclepias tuberosa*, is much favored by bumble bees. Aug-early Sept. Native, perennial, late to emerge in spring. Protect from slug damage when first sprouts appear in early June.



Borage, *Borago officinalis*, either blue or white, is eagerly visited by honey bees, bumble bees, sweat bees, and other small native bees. July-Aug. Annual, self-sows.



White meadowsweet, *Spiraea alba* var. *latifolia*, attracts bumble bees, sweat bees, and copious other insects. July-Sept. Native shrub, hardy and easy to grow. May already be in your area.



Summersweet, *Clethra alnifolia*, attracts many bees. Aug-Sept. Hardy native shrub.



Purple coneflower, *Echinacea purpurea*, has often been recommended for bee gardens, attracts bumble bees over a long season. Aug-Sept. Hardy perennial, easy to grow.



White wood aster, *Eurybia divaricata*, is one of many native asters that attracts Honey bees, bumble bees, and sweat bees. Sept. Hardy perennial, easy, spreads.



Greek oregano, *Origanum vulgare hirtum*, is popular with bumble bees and sweat bees. June-Aug. Hardy perennial, easy to grow, highly edible for people, too.

In cooperation with Dr. Francis A. Drummond, Dr. Lois Berg Stack, and Eric Venturini. Funding provided by the USDA and the University of Maine.