

NEW

State of Maine**Master Agreement****Effective Date:** 04/29/19**Expiration Date:** 04/30/20**Master Agreement Description:** Herbicides - Method 240 SL**Buyer Information**

William Allen

207-624-7871 ext. NULL WJE.Allen@maine.gov

Issuer Information

SHARON KRECHKIN

207-624-3038 ext. sharon.krechkin@maine.gov

Requestor Information

Sharon Krechkin

207-624-3038 ext. sharon.krechkin@maine.gov

Authorized Departments17A TRANSPORTATION
ALL**Vendor Information****Vendor Line #:** 1**Vendor ID**

VC1000093306

Vendor Name

NUTRIEN AG SOLUTIONS INC

Alias/DBA**Vendor Address Information**

18 LEGATE HILL RD

STERLING, MA 01564

US

Vendor Contact Information

RON LEMIN

207-944-6160 ext.

RONALD.LEMIN@NUTRIEN.COM

Commodity Information

Vendor Line #: 1

Vendor Name: NUTRIEN AG SOLUTIONS INC

Commodity Line #: 1

Commodity Code: 67590

Commodity Description: Herbicides - Method 240 SL

Commodity Specifications:

Commodity Extended Description: Method 240 SL

Quantity

0.00000

UOM

Unit Price

\$0.00

Delivery Days

3

Free on Board

Contract Amount

\$0.00

Service Start Date

Service End Date

Catalog Name

Method 240 SI

Discount

0.0000 %

Discount Start Date

04/29/19

Discount End Date

04/30/20

TERMS & CONDITIONS
MA 190426-162

COMMODITY ITEM: Herbicides - Method 240 SL

CONTRACT PERIOD: Through April 30, 2020. The State of Maine with vendor approval can opt to issue up to two (2) one (1) year extensions.

VENDOR CONTACT PERSON: The contact person will help consumers place orders, inquire about orders that have not been delivered, all shipping issues, quality issues, and any issues pertaining to this Master Agreement. All orders not submitted through a DO will be sent through the contractor's contact person. The contact person will be: **Ronald Lemin** Phone: **207-944-6160**
Email: Ronals.lemin@nutrien.com

EXTENSION OF CONTRACT: The Director of Procurement Services may, with the consent of the contractor extend the Contract period beyond the indicated expiration date.

CANCELLATION OF CONTRACT: The Division of Procurement Services reserves the right to cancel a contract with a thirty-day written notice OR cancel immediately if the contractor does not conform to terms and conditions and specifications of contract.

PRICES: Pricing will be FOB destination to all MaineDOT facilities. Prices are to remain firm for the duration of the contract. The State intends for this to mean that all goods shall be priced to include shipping charges, if any, to the State's desired location. The "FOB – Destination" shipping term is also intended to mean that the State shall not bear any responsibility for the goods in question until the State takes possession of them at the destination point of delivery.

DELIVERY: Product is required to be shipped to Topsham, West Gardiner, Dixfield, Eddington and Presque Isle. Other ship to points may be added in the future.

QUANTITIES: It is understood and agreed that the contract will cover the **actual quantities** required by State Agency over the length of the contract.

ORDERING PROCEDURE: Delivery orders (DO) will be created in AdvantageME for all orders over \$5000.00. If a DO is used, the DO will be e-mailed to the email address set up in AdvantageME by the Vendor as a .pdf file. Orders less than \$5000.00 can be ordered using a P-Card.

INVOICES and PAYMENT: The Contractor shall submit an itemized bill to the Department for materials following delivery for approval and payment. Invoices shall minimally include the following:
Contractor name, address & Contract Number 18P 190426-162, Invoice Date & Number

QUARTERLY REPORT: The Division of Procurement Services **requires a quarterly report of sales** be faxed to 207-287-6578 within 30 days of the end of each calendar quarter. It will be the responsibility of the vendor to produce a quarterly report. The report must include the dollar value of goods purchased, broken down by Department as well as the total dollar value of purchases made by all Departments.

PROCUREMENT CARD: State policy requires vendors to accept the State of Maine Procurement Card (P-Card) as a form of payment, with very rare exceptions. Your company is required to accept these cards. The pricing offered to the State of Maine shall be the final cost to the State of Maine regardless of payment method. No surcharge or other compensation will be allowed. The State of Maine reserves the right to reject your bid if you are unwilling to accept this condition.

ITEM DESCRIPTION	EXTENDED DESCRIPTION	UNIT OF MEASURE	LIST PRICE	DELIVERY DAYS
Herbicide, Metod 240 SL, Case	Case contains 2, 2.5 gallon containers	CASE	\$1,593.60	3



METHOD® 240SL HERBICIDE

Version 2.0 / USA
102000030323

1/8
Revision Date: 09/02/2015
Print Date: 11/18/2015

SECTION 1: IDENTIFICATION OF THE SUBSTANCE/MIXTURE AND OF THE COMPANY/UNDERTAKING

Product identifier

Trade name METHOD® 240SL HERBICIDE

Product code (UVP) 84117099

SDS Number 102000030323

EPA Registration No. 432-1565

Relevant identified uses of the substance or mixture and uses advised against

Use Herbicide

Restrictions on use See product label for restrictions.

Information on supplier

Supplier Bayer Environmental Science
2 T.W. Alexander Drive
Research Triangle PK, NC 27709
United States

Responsible Department Email: SDSINFO.BCS-NA@bayer.com

Emergency telephone no.

Emergency Telephone Number (24hr/ 7 days) 1-800-334-7577

Product Information Telephone Number 1-800-331-2867

SECTION 2: HAZARDS IDENTIFICATION

Classification in accordance with regulation HCS 29CFR §1910.1200

This material is not hazardous under the criteria of the Federal OSHA Hazard Communication Standard 29CFR 1910.1200.

Other hazards

No particular hazards known.

SECTION 3: COMPOSITION/INFORMATION ON INGREDIENTS

Hazardous Component Name	CAS-No.	Concentration % by weight
Aminocyclopyrachlor	858956-08-8	21.2



METHOD® 240SL HERBICIDE

Version 2.0 / USA
102000030323

2/8
Revision Date: 09/02/2015
Print Date: 11/18/2015

SECTION 4: FIRST AID MEASURES

Description of first aid measures

General advice	When possible, have the product container or label with you when calling a poison control center or doctor or going for treatment.
Inhalation	Move to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a physician or poison control center immediately.
Skin contact	Take off contaminated clothing and shoes immediately. Wash off immediately with plenty of water for at least 15 minutes. Call a physician or poison control center immediately.
Eye contact	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a physician or poison control center immediately.
Ingestion	Call a physician or poison control center immediately. Rinse out mouth and give water in small sips to drink. DO NOT induce vomiting unless directed to do so by a physician or poison control center. Never give anything by mouth to an unconscious person. Do not leave victim unattended.

Most important symptoms and effects, both acute and delayed

Symptoms	No symptoms known or expected.
Indication of any immediate medical attention and special treatment needed	
Treatment	Appropriate supportive and symptomatic treatment as indicated by the patient's condition is recommended. There is no specific antidote.

SECTION 5: FIREFIGHTING MEASURES

Extinguishing media

Suitable	Water spray, Foam, Dry chemical, Carbon dioxide (CO2)
Unsuitable	None known.

Advice for firefighters

Special protective equipment for fire-fighters	Firefighters should wear NIOSH approved self-contained breathing apparatus and full protective clothing.
---	--

Further information	Keep out of smoke. Fight fire from upwind position. Cool closed containers exposed to fire with water spray. Do not allow run-off from fire fighting to enter drains or water courses.
----------------------------	--

Flash point	> 100 °C
Autoignition temperature	no data available
Lower explosion limit	no data available
Upper explosion limit	no data available



METHOD® 240SL HERBICIDE

Version 2.0 / USA
102000030323

3/8
Revision Date: 09/02/2015
Print Date: 11/18/2015

Explosivity not applicable

SECTION 6: ACCIDENTAL RELEASE MEASURES

Personal precautions, protective equipment and emergency procedures

Precautions Keep unauthorized people away. Isolate hazard area. Avoid contact with spilled product or contaminated surfaces.

Methods and materials for containment and cleaning up

Methods for cleaning up Soak up with inert absorbent material (e.g. sand, silica gel, acid binder, universal binder, sawdust). Collect and transfer the product into a properly labelled and tightly closed container. Clean contaminated floors and objects thoroughly, observing environmental regulations.

Additional advice Do not allow to enter soil, waterways or waste water canal.

Reference to other sections Information regarding safe handling, see section 7.
Information regarding personal protective equipment, see section 8.
Information regarding waste disposal, see section 13.

SECTION 7: HANDLING AND STORAGE

Precautions for safe handling

Advice on safe handling Handle and open container in a manner as to prevent spillage. Use only in area provided with appropriate exhaust ventilation.

Hygiene measures Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, using the toilet or applying cosmetics.
Remove Personal Protective Equipment (PPE) immediately after handling this product. Before removing gloves clean them with soap and water. Remove soiled clothing immediately and clean thoroughly before using again. Wash thoroughly and put on clean clothing.

Conditions for safe storage, including any incompatibilities

Requirements for storage areas and containers Store in original container. Store in a cool, dry place and in such a manner as to prevent cross contamination with other crop protection products, fertilizers, food, and feed.

SECTION 8: EXPOSURE CONTROLS/PERSONAL PROTECTION

Control parameters

No control parameters known.

Exposure controls



METHOD® 240SL HERBICIDE

Version 2.0 / USA
102000030323

4/8
Revision Date: 09/02/2015
Print Date: 11/18/2015

Personal protective equipment

In normal use and handling conditions please refer to the label and/or leaflet. In all other cases the following recommendations would apply.

Respiratory protection	When respirators are required, select NIOSH approved equipment based on actual or potential airborne concentrations and in accordance with the appropriate regulatory standards and/or industry recommendations.
Hand protection	Chemical resistant nitrile rubber gloves
Eye protection	Chemical resistant goggles must be worn.
Skin and body protection	Wear long-sleeved shirt and long pants and shoes plus socks.
General protective measures	Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and warm/tepid water. Keep and wash PPE separately from other laundry.

SECTION 9. PHYSICAL AND CHEMICAL PROPERTIES

Appearance	brown
Physical State	liquid, clear
Odor	characteristic
Odour Threshold	no data available
pH	6.9 at 1 %
Vapor Pressure	no data available
Vapor Density (Air = 1)	no data available
Density	1.13 g/cm ³
Evaporation rate	no data available
Boiling Point	no data available
Melting / Freezing Point	no data available
Water solubility	soluble
Minimum Ignition Energy	not applicable
Decomposition temperature	not applicable
Partition coefficient: n-octanol/water	no data available
Viscosity	no data available
Flash point	> 100 °C
Autoignition temperature	no data available
Lower explosion limit	no data available



METHOD® 240SL HERBICIDE

Version 2.0 / USA
102000030323

5/8
Revision Date: 09/02/2015
Print Date: 11/18/2015

Upper explosion limit no data available
Explosivity not applicable

SECTION 10: STABILITY AND REACTIVITY

Reactivity

Thermal decomposition not applicable
Chemical stability Stable under recommended storage conditions.
Possibility of hazardous reactions No dangerous reaction known under conditions of normal use.
Conditions to avoid no data available
Incompatible materials no data available
Hazardous decomposition products No decomposition products expected under normal conditions of use.

SECTION 11: TOXICOLOGICAL INFORMATION

Immediate Effects
Eye Moderate eye irritation.

Information on toxicological effects

Acute oral toxicity LD50 (rat) > 5,000 mg/kg
Acute inhalation toxicity LC50 (rat) > 6.9 mg/l
Exposure time: 4 h
Acute dermal toxicity LD50 (rat) > 5,000 mg/kg
Skin irritation No skin irritation (rabbit)
Eye irritation Moderate eye irritation. (rabbit)
Sensitisation Non-sensitizing. (mouse)

ACGIH

None.

NTP

None.

IARC

None.



METHOD® 240SL HERBICIDE

Version 2.0 / USA
102000030323

6/8
Revision Date: 09/02/2015
Print Date: 11/18/2015

OSHA

None.

Further information

Only acute toxicity studies have been performed on the formulated product.

SECTION 12: ECOLOGICAL INFORMATION

Toxicity to fish	LC50 (Oncorhynchus mykiss (rainbow trout)) > 119 mg/l Exposure time: 96 h
Chronic toxicity to fish	Oncorhynchus mykiss (rainbow trout) NOEC: 11 mg/l Exposure time: 90 d
Toxicity to aquatic invertebrates	EC50 (Daphnia magna (Water flea)) 43 mg/l Exposure time: 96 h
Chronic toxicity to aquatic invertebrates	NOEC (Daphnia magna (Water flea)): 6 mg/l Exposure time: 21 d
Toxicity to aquatic plants	EC50 (Anabaena flos-aquae (cyanobacterium)) > 7.4 mg/l Exposure time: 72 h EC50 (Anabaena flos-aquae (cyanobacterium)) > 119 mg/l Exposure time: 96 h
Environmental precautions	Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate surface or ground water by cleaning equipment or disposal of wastes, including equipment wash water. Apply this product as specified on the label.

SECTION 13: DISPOSAL CONSIDERATIONS

Waste treatment methods

Product	Do not contaminate water, food, or feed by disposal. Dispose in accordance with all local, state/provincial and federal regulations. Follow advice on product label and/or leaflet.
Contaminated packaging	Do not re-use empty containers. Triple rinse containers.



METHOD® 240SL HERBICIDE

Version 2.0 / USA
102000030323

7/8
Revision Date: 09/02/2015
Print Date: 11/18/2015

Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or incineration, or if allowed by State and Local authorities, by burning.
If burned, stay out of smoke.
Follow advice on product label and/or leaflet.

RCRA Information

Characterization and proper disposal of this material as a special or hazardous waste is dependent upon Federal, State and local laws and are the user's responsibility. RCRA classification may apply.

SECTION 14: TRANSPORT INFORMATION

According to national and international transport regulations this material is not classified as dangerous goods / hazardous material.

SECTION 15: REGULATORY INFORMATION

EPA Registration No. 432-1565

US Federal Regulations

TSCA list

None.

US. Toxic Substances Control Act (TSCA) Section 12(b) Export Notification (40 CFR 707, Subpt D)

None.

SARA Title III - Section 302 - Notification and Information

None.

SARA Title III - Section 313 - Toxic Chemical Release Reporting

None.

US States Regulatory Reporting

CA Prop65

This product does not contain any substances known to the State of California to cause cancer.

This product does not contain any substances known to the State of California to cause reproductive harm.

US State Right-To-Know Ingredients

None.

Canadian Regulations

Canadian Domestic Substance List

None.

Environmental

CERCLA

None.

Clean Water Section 307 Priority Pollutants

None.

Safe Drinking Water Act Maximum Contaminant Levels



METHOD® 240SL HERBICIDE

Version 2.0 / USA
102000030323

8/8
Revision Date: 09/02/2015
Print Date: 11/18/2015

None.

EPA/FIFRA Information:

This chemical is a pesticide product registered by the Environmental Protection Agency and is subject to certain labeling requirements under federal pesticide law. These requirements differ from the classification criteria and hazard information required for safety data sheets, and for workplace labels of non-pesticide chemicals. Following is the hazard information required on the pesticide label:

Signal word: Caution!
Hazard statements: Moderate eye irritation.
Avoid contact with eyes.

SECTION 16: OTHER INFORMATION

Abbreviations and acronyms

49CFR	Code of Federal Regulations, Title 49
ACGIH	US. ACGIH Threshold Limit Values
CAS-Nr.	Chemical Abstracts Service number
EINECS	European inventory of existing commercial substances
ELINCS	European list of notified chemical substances
IARC	US. IARC Monographs on Occupational Exposures to Chemical Agents
IATA	International Air Transport Association
IMDG	International Maritime Dangerous Goods
N.O.S.	Not otherwise specified
NTP	US. National Toxicology Program (NTP) Report on Carcinogens
OECD	Organization for Economic Co-operation and Development
TDG	Transportation of Dangerous Goods
TWA	Time weighted average
UN	United Nations
WHO	World health organisation

NFPA 704 (National Fire Protection Association):

Health - 1 Flammability - 1 Instability - 0 Others - none

HMIS (Hazardous Materials Identification System, based on the Third Edition Ratings Guide)

Health - 1 Flammability - 1 Physical Hazard - 0 PPE -

0 = minimal hazard, 1 = slight hazard, 2 = moderate hazard, 3 = severe hazard, 4 = extreme hazard

Reason for Revision: New Safety Data Sheet.

Revision Date: 09/02/2015

This information is provided in good faith but without express or implied warranty. The customer assumes all responsibility for safety and use not in accordance with label instructions. The product names are registered trademarks of Bayer.

DO NOT USE PLANT MATERIAL TREATED WITH
METHOD® 240SL HERBICIDE FOR MULCH OR COMPOST



Method® 240SL

HERBICIDE

Soluble Liquid
For Non-Crop Use

ACTIVE INGREDIENT:	By Weight
Potassium salt of aminocyclopyrachlor	
Potassium salt of 6-amino-5-chloro-2-cyclopropyl-4-pyrimidinecarboxylic acid*	25%
OTHER INGREDIENTS:	75%
TOTAL:	100%

*Acid Equivalent: 6-Amino-5-chloro-2-cyclopropyl-4-pyrimidinecarboxylic acid
- 2 pounds acid per gallon or 21.2%

EPA REG. NO. 432-1565

Nonrefillable Container

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

See inside leaflet for complete First Aid
Instructions, Precautionary Statements, Directions
for Use and Storage and Disposal Instructions.

Net Contents
2.5 Gallons
84099295
84942561A 150820AV1

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers and loaders must wear:

Long-sleeved shirt and long pants. Shoes plus socks.

Applicators: After the product has been diluted in accordance with label directions for use, shirt, pants, socks, and shoes are sufficient Personal Protective Equipment (PPE).

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statement: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-334-7577 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of aminocyclopyrachlor from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

Ground Water Advisory

Aminocyclopyrachlor has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the watertable is shallow.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

METHOD® 240SL HERBICIDE must be used only in accordance with directions on this label or in separately published BAYER CROPS-SCIENCE LP directions.

BAYER CROPS-SCIENCE LP will not be responsible for losses or damages resulting from the use of this product in any manner not specifically directed by BAYER CROPS-SCIENCE LP. User assumes all risks associated with such non-directed use.

PRODUCT INFORMATION

METHOD 240SL HERBICIDE is a soluble liquid that is mixed in water and applied as a spray. METHOD 240SL HERBICIDE may be applied by aerial or ground equipment for control of broadleaf weeds and woody species, including many terrestrial and riparian invasive and noxious weeds. METHOD 240SL HERBICIDE is registered for general weed and brush control on private, public, and military lands as follows: uncultivated non-agricultural areas (such as airports, highway, railroad and utility rights-of-way, sewage disposal areas, etc.); uncultivated agricultural areas - non-crop producing (such as farmyards, fuel storage areas, fence rows, non-irrigation ditchbanks, barrier strips, etc.); industrial sites - outdoor (such as lumberyards, pipeline and tank farms, etc.); and natural areas (such as wildlife management areas, wildlife openings, and wildlife habitats). METHOD 240SL HERBICIDE may be used for the release or restoration of native perennial grasses and in established industrial turf grasses.

This product may be applied to terrestrial non-crop sites and unimproved turf sites that contain areas of temporary surface water, caused by collection of water in equipment ruts or in other depressions created by management activities. It is permissible to treat intermittently flooded low lying sites, seasonally dry flood plains, and transitional areas between upland and lowland sites when no water is present. It is also permissible to treat marshes, swamps, and bogs after water has receded, as well as seasonally dry flood deltas. METHOD 240SL HERBICIDE may be applied up to the water's edge. Do not apply directly to water.

METHOD 240SL HERBICIDE provides preemergence and/or postemergence control of the broadleaf weeds, vines, and brush species listed in the weeds controlled section of the label. For perennial species on the label, a postemergence application should be used. For best postemergence performance, an MSO type adjuvant should be included to the spray solution. Excessive wetting of the target plant is not necessary but good spray coverage of the target plant is needed for best results.

METHOD 240SL HERBICIDE is non-corrosive to spray equipment.

Do not apply more than 18 fluid ounces per acre per year.

BIOLOGICAL ACTIVITY

METHOD 240SL HERBICIDE is quickly taken up by the leaves, stems and roots of plants. The effects of METHOD 240SL HERBICIDE may be seen on plants from within a few hours to a few days. The most noticeable symptom is a bending and twisting of stems and leaves. Other advanced symptoms include severe necrosis, stem thickening, growth stunting, leaf crinkling, calloused stems and leaf veins, leaf-cupping, and enlarged roots. Death of treated broadleaf plants may require several more weeks and up to several months for some woody plant species. METHOD 240SL HERBICIDE is rain-fast at 1 hour after application.

IMPORTANT RESTRICTIONS

- Do not apply this product in areas where the roots of desirable trees and/or shrubs may extend unless injury or loss can be tolerated. Root zone areas of desirable trees or vegetation are affected by local conditions and can extend well beyond the tree canopy.

- Do not apply this product if site-specific characteristics and conditions exist that could contribute to movement and unintended root zone exposure to desirable trees or vegetation, unless injury or loss can be tolerated.
- Do not make applications when circumstances favor movement from treatment site.
- Do not apply METHOD 240SL HERBICIDE to roadsides or other non-crop areas during periods of intense rainfall or where prevailing soils are either saturated with water or of a type through which rainfall will not readily penetrate as this may result in off-site movement.
- Do not apply or otherwise permit this product or sprays containing this product to come into contact with any non-target crop or desirable vegetation.
- Do not apply in or on dry or water containing irrigation ditches or canals including their outer banks.
- Do not apply through any type of irrigation system.
- Do not contaminate water intended for irrigation. To avoid injury to crops or other desirable vegetation, do not treat or allow spray drift or run-off to fall onto banks or bottoms of irrigation ditches, either dry or containing water, or other channels that carry water that may be used for irrigation purposes.
- Treatment of powdery dry soil and light sandy soils, when there is little likelihood of rainfall soon after treatment, may result in off target movement and possible damage to susceptible crops and desirable vegetation, when soil particles are moved by wind or water. Injury to crops or desirable vegetation may result if treated soil is washed, blown or moved onto land used to produce crops or land containing desirable vegetation. Do not apply METHOD 240SL HERBICIDE when these conditions are identified and powdery dry soil or light or sandy soils are known to be prevalent in the area to be treated.
- Do not apply when the soil is frozen or covered with snow.
- Do not use on lawns, walks, driveways, tennis courts, or similar areas.
- Do not apply more than 18 fluid ounces (0.28 pound ae) per acre per year.
- Do not graze or feed forage, hay, or straw from treated areas to livestock.
- Do not use plant material treated with this product for mulch or compost.
- If non-crop sites treated with METHOD 240SL HERBICIDE are to be converted to a food, feed, or fiber agricultural crop, or to a horticultural crop, do not plant the treated sites for at least one year after the METHOD 240SL HERBICIDE application. A field bioassay must then be completed before planting the desired crop.
- Not for sale, sale into, distribution, and/or use in Nassau and Suffolk counties of New York State.

IMPORTANT PRECAUTIONS

- Certain species may, in particular, be sensitive to low levels of METHOD 240SL HERBICIDE including but not limited to conifers (such as Douglas fir, Norway spruce, ponderosa pine and white pine), deciduous trees (such as aspen, Chinese tallow, cottonwood, honey locust, magnolia, poplar species, redbud, silver maple, and willow species), and ornamental shrubs (such as arborvitae, burning bush, crape myrtle, forsythia, hydrangea, ice plant, magnolia, purple plum and yew).
- Injury or loss of desirable trees or vegetation may result if METHOD 240SL HERBICIDE is applied on or near desirable trees or vegetation, on areas where their roots extend, or in locations where the treated soil may be washed or moved into contact with their roots. Consider site-specific characteristics and conditions that could contribute to unintended root zone exposure to desirable trees or vegetation. Root zone areas of desirable trees or vegetation are affected by local conditions and can extend beyond the tree canopy. If further information is needed regarding root zone area, consult appropriate state extension service, professional consultant, or other qualified authority.
- Injury to or loss of desirable trees or vegetation may result if equipment is drained or flushed on or near these trees or vegetation or on areas where their roots may extend or in locations where the chemical may be washed or moved into contact with their roots.
- In non-crop areas adjacent to desirable vegetation, avoid overlapping spray applications and shut off spray to the spray boom while starting, turning, slowing, or stopping to avoid injury to desirable vegetation.
- Applications made where runoff water flows onto agricultural land may injure or kill crops such as but not limited to sugar beets, potatoes, tomatoes, tobacco, soybeans, field beans, alfalfa, grapes, peaches, almonds, and vegetables.
- Applications should be made only when there is little or no hazard from spray drift. Very small quantities of spray, which may not be visible, may seriously injure susceptible plants.
- Exposure to METHOD 240SL HERBICIDE may injure or kill most crops and may injure or kill desirable vegetation. Injury may be more severe when the crops or desirable vegetation are irrigated.
- Caution is advised when using this product in areas where loss of desirable conifer or deciduous trees and/or shrubs, as well as other broadleaf plants, including but not limited to legumes and wild flowers, cannot be tolerated. Without prior experience, it is necessary that small areas containing these plants be tested for tolerance to METHOD 240SL HERBICIDE and its soil residues before any large scale spraying occurs.
- Low rates of METHOD 240SL HERBICIDE can kill or severely injure most crops. Following a METHOD 240SL HERBICIDE application, the use of spray equipment to apply other pesticides to crops on which METHOD 240SL HERBICIDE is not registered may result in their damage. The most effective way to reduce this crop damage potential is to use dedicated mixing and application equipment.
- Leave treated soil undisturbed to reduce the potential for METHOD 240SL HERBICIDE movement by soil erosion due to wind or water.
- In the case of suspected off-site movement of METHOD 240SL HERBICIDE to cropland, soil samples should be quantitatively analyzed for METHOD 240SL HERBICIDE, or any other herbicide which could be having an adverse effect on the crop, in addition to conducting the field bioassay.
- METHOD 240SL HERBICIDE may suppress or severely injure certain established grasses, such as some brome grass and wheat-grass species, especially when the grass plants are stressed by adverse environmental conditions. Areas that contain these grass plants should recover as environmental conditions for good grass growth occur.

FIELD BIOASSAY

To conduct a field bioassay, grow to maturity test strips of the crop you plan to grow the following year. The test strips should cross the entire field including knolls and low areas. Crop response to the field bioassay will indicate whether or not to plant the crops grown in the test strips. If no crop injury (such as, poor germination, stunting, or chlorosis, malformation, or necrosis of leaves) or yield loss is evident from the crops grown in the test strips, the intended rotational crop may be planted. If herbicide symptoms or yield loss is observed, do not plant the crop.

TANK MIXTURES

METHOD 240SL Herbicide may be tank mixed with other herbicides which are registered for the same use sites, methods of application, and timings as specified on this product label. Refer to the tank mix product label for any additional instructions or use restrictions. In addition, a spray adjuvant may be mixed with METHOD 240SL HERBICIDE when making postemergence applications. Refer to the adjuvant label for additional instructions or use restrictions.

ADJUVANTS

Methylated Seed Oils and Vegetable Oils: A methylated seed oil (MSO) or vegetable oil based adjuvant may provide increased leaf absorption

of METHOD 240SL HERBICIDE. Include the MSO or vegetable oil adjuvant at 1% w/v (1 gallon per 100 gallons of spray solution).

Non-ionic Surfactants: Use a non-ionic surfactant at a minimum rate of 0.25% v/v (1 quart surfactant per 100 gallons of spray solution). Surfactant products must contain at least 70% non-ionic surfactant with a hydrophilic/lipophilic balance (HLB) of 12 to 17.

Invert Emulsions: METHOD 240SL HERBICIDE may be applied as an invert emulsion. The spray solution results in an invert (water- in-oil) spray emulsion designed to minimize spray drift and spray run-off, resulting in more herbicide deposited on the target foliage. The spray emulsion may be formed in a single tank (batch mixing) or injected (in-line mixing). Consult the invert chemical label for proper mixing directions.

INVASIVE SPECIES MANAGEMENT

This product may be used on public, private, and tribal lands to treat certain weed species infestations that have been determined to be invasive, consistent with the Federal Interagency Committee for the Management of Noxious and Exotic Weeds (FICMNEW) National Early Detection and Rapid Response (EDRR) System for invasive plants. Effective EDRR systems address invasions by eradicating the invader where possible, and controlling them when the invasive species is too established to be feasibly eradicated. Once an EDRR assessment has been completed and action is recommended, a Rapid Response needs to be taken to quickly contain, deny reproduction, and, if possible, eliminate the invader. Consult your appropriate state extension service, forest service, or regional multidisciplinary invasive species management coordination team to determine the appropriate Rapid Response provisions and allowed treatments in your area.

RESISTANCE

When herbicides that affect the same biological site of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field.

Adequate control of these resistant weed biotypes cannot be expected. If weed control is unsatisfactory, it may be necessary to retreat the problem area using a product affecting a different site of action. To better manage herbicide resistance through delaying the proliferation and possible dominance of herbicide resistant weed biotypes, it may be necessary to change cultural practices within and between crop seasons such as using a combination of tillage, retreatment, tank-mix partners and/or sequential herbicide applications that have a different site of action. Weed escapes that are allowed to go to seed will promote the spread of resistant biotypes. It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes. Consult your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative for specific alternative cultural practices or herbicide recommendations available in your area.

INTEGRATED PEST MANAGEMENT

This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

NON-AGRICULTURAL USES

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Terrestrial non-crop weed control is not within the scope of the Worker Protection Standard. See the Product Information section of this label for a description of noncrop sites.

Do not enter terrestrial/non-crop treated areas without protective clothing until sprays have dried.

PRODUCT INFORMATION FOR NON-AGRICULTURAL USES

METHOD 240SL HERBICIDE is a soluble liquid that is mixed in water and applied as a spray. METHOD 240SL HERBICIDE may be applied by aerial or ground equipment for control of broadleaf weeds and woody species, including many terrestrial and riparian invasive and noxious weeds. METHOD 240SL HERBICIDE is registered for general weed and brush control on private, public, and military lands as follows: uncultivated non-agricultural areas (such as airports, highway, railroad and utility rights-of-way, sewage disposal areas, etc.); uncultivated agricultural areas - noncrop producing (such as farmyards, fuel storage areas, fence rows, non-irrigation ditch banks, barrier strips, etc.); industrial sites - outdoor (such as lumberyards, pipeline and tank farms, etc.); and natural areas (such as wildlife management areas, wildlife openings, and wildlife habitats).

METHOD 240SL HERBICIDE may be used for the establishment or release of native grasses and for weed control in established, unimproved grass turf.

Apply METHOD 240SL HERBICIDE preemergence or early postemergence when broadleaf weeds are actively germinating or growing. METHOD 240SL HERBICIDE can provide long-term control of susceptible weeds. The length of control is dependent upon the application rate, condition and growth stage of target weeds, environmental conditions at and following application, and the density and vigor of competing desirable vegetation. Best results for long term weed control occur where grasses and other vegetation is allowed to recover from adverse environmental conditions and compete with susceptible weeds.

Weeds hardened off by cold weather or drought stress may not be controlled.

METHOD 240SL HERBICIDE may be applied broadcast using ground spray equipment, fixed-wing aircraft, or by helicopter. When applying by fixed-wing aircraft or helicopter, follow directions under the Aerial Applications section of this label; otherwise refer to the section on Ground Applications when using surface equipment.

METHOD 240SL HERBICIDE may also be applied using low and high volume ground spray equipment.

APPLICATION INFORMATION

AERIAL APPLICATIONS

When applying by air, apply only using nozzles which will deliver coarse or greater (VMD >350 microns) droplets as defined by ASABE S572 standard. Do not release spray at a height greater than 10 feet above the ground or canopy unless a greater height is required for aircraft safety. Do not apply when wind speed is greater than 10 mph. Do not apply during a temperature inversion.

For aerial applications near susceptible crops or other desirable plants, use a drift control additive as recommended by the manufacturer, or apply through a "Microfoil" or "Thru-Valve" boom, or use an equivalent drift control system. Thickened sprays prepared by using high viscosity invert systems, or other drift control systems, may be utilized if drift control is comparable to that obtained with drift control additives or the "Thru-Valve" boom. If a spray thickening agent is used, follow all recommendations and precautions on the product label. Do not use a thickening agent with the "Microfoil" boom or other systems that cannot accommodate thick sprays.

METHOD 240SL HERBICIDE may be applied by either fixed-wing aircraft or helicopter spray equipment. Fixedwing aircraft and helicopters can be used to apply METHOD 240SL HERBICIDE; however, do not make applications by fixed-wing aircraft unless appropriate buffer zones can be maintained to prevent spray drift out of the target area or, when treating open tracts of land, spray drift

as a result of fixed wing aircraft application can be tolerated. The application volume required will vary with the height and density of the brush and the application equipment used. Generally, aerial applications will require 15 to 25 gallons of spray solution per acre. Regardless of the application volume or spray equipment used, thorough coverage of the foliage is necessary to optimize control results.

All precautions and restrictions should be taken to minimize or eliminate spray drift.

GROUND APPLICATIONS

When applying by ground, apply only using nozzles which will deliver coarse or greater (VMD >350 microns) droplets as defined by ASABE S572 standard. Do not apply with a nozzle height greater than 4 feet above the ground or canopy unless necessitated by the application equipment. Apply with the spray boom or nozzle height as low as possible. Do not apply when wind speed is greater than 10 mph. Do not apply during a temperature inversion.

For ground applications, keep the spray boom as low as possible; apply 10 gallons or more of spray per acre; use spray pressures no greater than are required to obtain adequate plant coverage; use large-droplet producing nozzle tips; use drift control additives; use shielded-sprayers or other drift control systems; and/or spray when wind velocity is low.

LOW-VOLUME FOLIAR APPLICATION

For low-volume applications, see Table 1 for use rate and mixing guidelines. The spray concentration of METHOD 240SL HERBICIDE should be adjusted according to the spray volume per acre and the size and plant density of the target brush species. For best results, include an MSO adjuvant at the rate of 1% v/v. Good plant coverage is necessary for best results. Use spray nozzles and pressure that will aid the proper deposition of the spray solution. Apply in sufficient spray volume to help provide uniform spray distribution of spray particles over the area to be treated and to avoid spray drift. Generally, low volume ground applications will require 20 to 50 gallons per acre and ultra-low volume ground application will require 10 to 20 gallons of spray solution per acre. The use of an even flat fan tip with a spray angle of 40 degrees or less will aid in proper spray deposition. Some recommended tip sizes include 4004E or 1504E. For cone or straight stream nozzle patterns, the adjustable cone nozzles, such as the 5500 X3 or the 5500 X4 may be used. Use the higher concentration rates for hard to control brush species. Do not apply more than 18 fluid ounces of METHOD 240SL HERBICIDE per acre per year. Note: Add a spray pattern indicator, if desired, at the recommended label rates.

HIGH VOLUME FOLIAR APPLICATION

High volume applications may be applied at rates equivalent to broadcast rates up to 18 fluid ounces per acre per year. Where a rate range is indicated for the brush species, use the higher rate for high density brush sites. For best results, use MSO adjuvant at the rate of 1% V/V to the spray solution.

When making broadcast applications, apply near the tops of the brush plants in a light drizzle pattern. The spray solution should reach the crown of the plants and trickle down into the canopy. Use sufficient spray volume to thoroughly and uniformly wet foliage and stems but don't over apply causing excessive run-off. Generally, high volume ground applications will require 100 to 400 gallons per acre. Do not apply more than 18 fluid ounces per broadcast acre per year.

Table 1: METHOD 240SL HERBICIDE Use Rate and Mixing Guide

Total Spray Volume (gallons per acre)	Rate of METHOD 240SL HERBICIDE 8 fluid ounces/acre [fluid ounces/ 100 gallons of spray]*	Rate of METHOD 240SL HERBICIDE 12 fluid ounces/acre [fluid ounces/ 100 gallons of spray]*	Rate of METHOD 240SL HERBICIDE 16 fluid ounces/acre [fluid ounces/ 100 gallons of spray]*
400	2	3	4
300	2.7	4	5.3
200	4	6	8
100	8	12	16
50	16	24	32
40	20	30	40
30	26.7	40	53.3
20	40	60	80
10	80	120	160

* Do not exceed the maximum use rate of 18 fluid ounces product broadcast per acre per year.

SPOT APPLICATION

Spot applications may be applied at rates equivalent to the broadcast application rate up to a maximum of 18 fluid ounces per acre per year. Use sufficient spray volume to thoroughly and uniformly wet target weed or brush foliage. Use of a high quality MSO adjuvant may be added to the spray mixture as recommended by the adjuvant manufacturer. Repeat applications may be made, but the total amount of METHOD 240SL HERBICIDE must not exceed 18 fluid ounces per year. To prevent misapplication, spot applications should be applied with either a calibrated boom sprayer, a boom-less sprayer, or a hand-held or backpack sprayer.

Do not apply more than 18 fluid ounces product per broadcast acre per year as a result of broadcast, spot, or repeat applications. Application rates in Table 2 are based on treating an area of 1000 square feet (sq ft). Mix METHOD 240SL HERBICIDE in 0.3 to 3 gallons of water, depending on the spray volume necessary to treat 1000 sq ft. A spray volume of 0.3 to 3 gallons per 1000 sq ft is equivalent to 13 to 130 gallons per acre.

Table 2. Spot spray use rates

Amount of METHOD 240SL HERBICIDE per 1000 square feet to Equal a Broadcast Rate		
Broadcast Rate (fl ounces/acre)	METHOD 240SL HERBICIDE needed per 1000 sq ft	
	(fl ounces)	(mls)
8	0.18	5.3
12	0.27	8
16	0.37	11
18	0.42	12.4

INVERT EMULSION APPLICATIONS

METHOD 240SL HERBICIDE can be applied as an invert emulsion (water in oil). This can be done in a batch mixing (single tank) or inline-mixing (injected) process. Follow the directions on the invert chemical guide.

CUT STUMP AND STEM TREATMENTS

Make a dilute solution by mixing 5 to 10 gallons of METHOD 240SL HERBICIDE in enough basal oil to make 100 gallons of spray mixture. Apply with a knapsack or backpack sprayer using low pressure and solid cone or flat fan nozzles. Spray the stump cut surface and thoroughly wet the cambium layer next to the bark, also treat the sides of the stump and the root collar area. On larger trees, treat only the outer 2-3 inches of the stump. On trees 3 inches or less in diameter treat the entire cut surface. Apply anytime except when snow or water prevents treating to the ground line of the stump. Moisture stress may affect optimum control.

BASAL BARK TREATMENTS

Make a dilute solution by mixing 10 to 20 gallons of METHOD 240SL HERBICIDE in enough basal oil to make 100 gallons of spray mixture. Apply with a knapsack or backpack sprayer using low pressure and solid cone or flat fan nozzles. Make applications to susceptible brush or tree species with stems less than 6 inches in basal diameter. Thoroughly wet the lower 12 to 18 inches of the trunk or stem (from ground line). Treat until run-off at the ground line is noticeable. Brush or trees with old or rough bark will require more spray solution than smooth young bark. Applications can be made anytime of the year except when snow or water prevents treating to the ground line of the brush or tree trunk.

CUT STUBBLE TREATMENTS

For the prevention of re-sprouting, after hand cutting or mechanical mowing of susceptible brush species along rights-of-way and other non-crop sites, apply a broadcast application of METHOD 240SL HERBICIDE at 18 fluid ounces product per acre. Apply in a minimum of 20 gallons of water per acre. Make applications soon after cutting. The addition of a penetrating agent at 5% V/V or more can aid in uptake through the bark or exposed roots of the cut brush. For best results, make applications before or during periods of active root growth. Do not apply when the soil is frozen or covered by standing water or snow.

SPECIFIC USE DIRECTIONS

BAREGROUND

METHOD 240SL HERBICIDE may be used in non-crop sites for bareground (total vegetation control) weed control. Preemergence or postemergence applications of METHOD 240SL HERBICIDE provide control of many annual and perennial broadleaf weeds. Apply at up to 18 fluid ounces product per acre in tank mixes with other products registered for use on bareground sites. Consult the manufacturer's labels for specific rates, weeds controlled and use restrictions.

Make a thorough and uniform application with calibrated spray equipment per label directions. Apply at any time of the year. Use the higher rates of METHOD 240SL HERBICIDE for fall applications and in previously untreated areas or areas with high weed infestations. For postemergence applications always include a spray adjuvant. For faster brown-out or burn down results, add glyphosate or similar products to the tank. For added residual weed control, or to broaden the weed control spectrum, tank mix with other residual products registered for use on bareground sites. The level and length of control will depend on the herbicide rate applied, amount of rainfall, soil texture, environmental and applications conditions.

UNIMPROVED TURF GRASS

METHOD 240SL HERBICIDE may be used in non-crop industrial sites, such as utility rights-of-way and roadsides, for general weed control in established industrial turf grasses. Apply METHOD 240SL HERBICIDE at 2.0 to 4.0 fluid ounces product per acre. Treatments made prior to the full green-up stage may delay green-up. Apply METHOD 240SL HERBICIDE by ground equipment only. Use a minimum of 10 gallons of water per acre. The addition of an MSO adjuvant may increase the potential for turf grass injury. Important: Temporary chlorosis (yellowing), reddening, stunting, droopy or twisted grass leaves, and seed head suppression may occur.

Do not apply in the first growing season of any grass. Do not apply METHOD 240SL HERBICIDE to grass under stress from disease, insects, drought, or other environmental causes.

NON-CROPLAND RESTORATION

METHOD 240SL HERBICIDE is labeled for the control of broadleaf weeds and brush, listed in the weeds controlled section, in unimproved industrial turf, on roadsides, airports, industrial sites, or on other similar non-crop sites in order to establish or release desirable, introduced or native perennial grass species for site stabilization.

To maximize and extend the weed and brush control provided by METHOD 240SL HERBICIDE, it is critical that other vegetation management practices, including mowing, fertilization, etc., be incorporated into the restoration program to help extend or build on the weed control benefits and promote the growth of introduced or established grasses and/or desirable plants or plant communities. During the season of establishment, METHOD 240SL HERBICIDE must only be applied after introduced or native perennial grasses are well established. The grass must have a good secondary root system and show good vigor.

METHOD 240SL HERBICIDE may suppress certain established grasses especially when the grass plants are stressed by adverse environmental conditions. Temporary reddening, stunting, droopy or twisted leaves may occur. Do not apply METHOD 240SL HERBICIDE to grass under stress from disease, insects, drought, or other environmental causes.

Apply METHOD 240SL HERBICIDE at 2.0 to 4.0 fluid ounces product per acre in the fall, before the soil freezes, or in the spring after the soil thaws. When applied at lower rates, METHOD 240SL HERBICIDE provides short-term control of weeds listed; when applied at higher rates, weed control spectrum is broadened and extended.

Do not apply when the soil is frozen.

WEEDS CONTROLLED

Use the higher spray volumes and herbicide rates for heavy weed and brush infestations, hard to control species, and tall brush or dense hardwood canopies. Do not apply more than 18 fluid ounces product broadcast per acre per year.

BROADLEAF WEEDS		Rate (fluid ounces per acre)
Clover, bush	<i>Lespedeza</i> sp.	4 to 8
Clover, Dutch (white)	<i>Trifolium repens</i>	
Dandelion, common	<i>Taraxacum officinale</i>	
Ironweed, tall	<i>Vernonia gigantea</i>	
Lespedeza, serotia	<i>Lespedeza cuneata</i>	
Lettuce, prickly	<i>Lactuca scariola</i>	
Mullein, turkey	<i>Croton salignus</i>	
Ragweed, western	<i>Ambrosia psilostachya</i>	
Sowthistle, common	<i>Sonchus oleraceus</i>	
Starthistle, yellow	<i>Centaurea solstitialis</i>	
Hawkweed, orange	<i>Hieracium aurantiacum</i>	8 to 18
Knopweed, diffuse	<i>Centaurea diffusa</i>	
Knopweed, Russian	<i>Centaurea repans</i>	
Knopweed, spotted	<i>Centaurea biebersteinii</i>	
Kochia (Up to 6 inches) ¹	<i>Kochia scoparia</i>	
Locust, honey	<i>Gleditsia triacanthos</i>	
Marestail/horseweed	<i>Conyza canadensis</i>	
Ragweed, common	<i>Ambrosia artemisiifolia</i>	
Spurge, leafy	<i>Euphorbia esula</i>	
Thistle, Canada	<i>Cirsium arvense</i>	
Thistle, cotton	<i>Onopordum acanthium</i>	
Thistle, musk	<i>Carduus nutans</i>	
Thistle, Russian	<i>Salsola iberica</i>	
Toadflax, dalmatian	<i>Linaria dalmatica</i>	
Plantain	<i>Plantago</i> spp.	10 to 18
Aster, white	<i>Aster pilosus</i>	12 to 18
Bindweed, field	<i>Convolvulus arvensis</i>	
Cinquefoil, sulfur	<i>Potentilla recta</i>	
Goldenrod, Canada	<i>Solidago canadensis</i>	
Hemlock, poison	<i>Conium maculatum</i>	
Honeysuckle, Japanese	<i>Lonicera japonica</i>	
Poison-ivy, eastern	<i>Toxicodendron radicans</i>	
Teasel	<i>Dipsacus fulvonum</i>	
Yarrow, common	<i>Achillea millefolium</i>	
BRUSH		Rate (fluid ounces per acre)
Ash (Green, White)	<i>Fraxinus</i> sp.	10 to 18
Catalpa	<i>Catalpa speciosa</i>	
Cottonwood	<i>Populus deltoides</i>	
Dewberry	<i>Rubus trivialis</i>	
Elder, box	<i>Acer negundo</i>	
Elm	<i>Ulmus americana</i>	
Hackberry, common	<i>Celtis occidentalis</i>	
Locust, black	<i>Robinia pseudoacacia</i>	
Maple, red	<i>Acer rubrum</i>	
Maple, silver	<i>Acer sacharinum</i>	
Poplar, yellow	<i>Liriodendron tulipifera</i>	
Sugarberry	<i>Celtis laevigata</i>	
Sumac	<i>Rhus</i> sp.	
Sycamore	<i>Acer pseudoplatanus</i>	
Tupelo, black	<i>Nyssa sylvatica</i>	
Willow, weeping	<i>Salix alba</i>	
Wild grape	<i>Vitis rotundifolia</i>	
Oak, northern red	<i>Quercus borealis</i>	16
Pine, Virginia ²	<i>Pinus virginia</i>	
Sassafras	<i>Sassafras albidum</i>	
Huisache	<i>Acacia farnesiana</i>	18
Mesquite	<i>Prosopis juliflora</i>	

¹-See specific weed directions.

²-Suppression: a visual reduction in weed competition (reduced population or vigor) as compared to an untreated area.

Specific Weed Directions:

Kochia: For non-selective applications, tankmixing glyphosate with Method@ 240 SL HERBICIDE may improve control under dry conditions.

SPRAY EQUIPMENT

Be sure the sprayer is calibrated before use. Use a sufficient volume of water that will deliver a uniform spray pattern and coverage of the target brush or weeds.

The selected sprayer should be equipped with an agitation system to help keep METHOD 240SL HERBICIDE suspended in the spray tank.

Note: Low rates of METHOD 240SL HERBICIDE can kill or severely injure most crops. Following an METHOD 240SL HERBICIDE application, the use of spray equipment to apply other pesticides to crops on which METHOD 240SL HERBICIDE is not registered may result in their damage.

The most effective way to reduce this crop damage potential is to use dedicated mixing and application equipment.

MIXING INSTRUCTIONS

1. Fill the tank 1/3 to 1/2 full of water.
2. While agitating, add the required amount of METHOD 240SL HERBICIDE.
3. Continue agitation until the METHOD 240SL HERBICIDE is fully dispersed, at least 5 minutes.
4. Once the METHOD 240SL HERBICIDE is fully dispersed, maintain agitation and continue filling tank with water. METHOD 240SL HERBICIDE should be thoroughly mixed with water before adding any other material.
5. As the tank is filling, add tank mix partners (if desired) and then add the necessary volume of spray adjuvants. Always add spray adjuvants last.
6. If the mixture is not continuously agitated, settling will occur. If settling occurs, thoroughly re-agitate before using.
7. Apply METHOD 240SL HERBICIDE spray mixture within 24 hours of mixing to avoid product degradation.
8. If METHOD 240SL HERBICIDE and a tank mix partner are to be applied in multiple loads, pre-slurry METHOD 240SL HERBICIDE in clean water prior to adding it to the tank. This will prevent the tank mix partner from interfering with the dissolution of the METHOD 240SL HERBICIDE.

SPRAYER CLEANUP

The spray equipment must be cleaned before METHOD 240SL HERBICIDE is sprayed. Follow the cleanup procedures specified on the labels of the previously applied products.

AT THE END OF THE DAY

It is recommended that, during periods when multiple loads of METHOD 240SL HERBICIDE are applied, at the end of each day of spraying the interior of the tank should be rinsed with fresh water and then partially filled and the boom and hoses flushed. This will prevent the buildup of dried pesticide deposits which can accumulate in the application equipment.

1. Empty the tank and drain the sump completely.
2. Spray the tank walls with clean water using a minimum volume of 10% of the tank volume. Circulate the water through the lines, including all by-pass lines, for at least two minutes. Flush the boom well and empty the sprayer. Completely drain the sump.
3. Repeat step 2.
4. Remove the nozzles and screens and clean separately in a bucket containing water. The rinsate solution may be applied to the non-crop sites listed on this label. Do not exceed the maximum labeled use rate. If cleaners are used, consult the cleaner label for rinsate disposal instructions. If no instructions are given, dispose of the rinsate on site or at an approved waste disposal facility.

Notes:

1. Always start with a clean spray tank.
2. Steam-cleaning aerial spray tanks is recommended to facilitate the removal of any caked deposits.
3. When METHOD 240SL HERBICIDE is tank mixed with other pesticides, all cleanout procedures for each product should be examined, and the most rigorous procedure should be followed.
4. In addition to this cleanout procedure, all pre-cleanout guidelines on subsequently applied products should be followed as per the individual labels.
5. Low rates of METHOD 240SL HERBICIDE can kill or severely injure most crops. Following a METHOD 240SL HERBICIDE application, the use of spray equipment to apply other pesticides to crops on which METHOD 240SL HERBICIDE or its active ingredients are not registered may result in their damage. The most effective way to reduce this crop damage potential is to use dedicated mixing and application equipment.

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

IMPORTANCE OF DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS! See Wind, Temperature and Humidity, and Temperature Inversions sections of this label.

CONTROLLING DROPLET SIZE - GENERAL TECHNIQUES

- Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.
- Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

CONTROLLING DROPLET SIZE - AIRCRAFT

- Number of Nozzles - Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- Nozzle Orientation - Orienting nozzles so that the spray is emitted backwards, parallel to the air stream will produce larger droplets than other orientations.
- Nozzle Type - Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.
- Boom Length - The boom length should not exceed 3/4 of the wing or rotor length - longer booms increase drift potential.
- Application Height - Application more than 10 ft above the canopy increases the potential for spray drift.

BOOM HEIGHT

Setting the boom at the lowest labeled height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom should remain level with the crop and have minimal bounce.

WIND

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. AVOID GUSTY OR WINDLESS CONDITIONS. Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

SURFACE TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind.

They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

SENSITIVE AREAS

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

DRIFT CONTROL ADDITIVES

Drift control additives may be used with all spray equipment with the exception of controlled droplet applicators. When a drift control additive is used, read and carefully observe cautionary statements and all other information on the label. It is recommended that drift control additives be certified by the Chemical Producers and Distributors Association (CPDA).

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store product in original container only. Store in a cool, dry place.

Pesticide Disposal: Waste resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

Refer to the Net Contents section of this product's labeling for the applicable "Nonrefillable Container" or "Refillable Container" designation.

Nonrefillable Rigid Plastic and Metal Containers (Capacity Equal to or Less Than 5 Gallons):

Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Rigid Plastic and Metal Containers (Capacity Greater Than 5 Gallons): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Rigid Plastic and Metal Containers, e.g., Intermediate Bulk Containers (IBC) (Size or Shape Too Large to be Tipped, Rolled or Turned Upside Down): Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying the contents from this container into application equipment or mix tank and before final disposal using the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour, or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

All Refillable Containers: Refillable container. Refilling Container: Refill this container with METHOD 240SL HERBICIDE containing aminocyclopyrachlor potassium salt only. Do not reuse this container for any other purpose. Cleaning before refilling is the responsibility of the refiller. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn out threads and closure devices. If damage is found, do not use container; contact BAYER CROPSCIENCE LP at the number below for instructions. Check for leaks after refilling and before transporting. If leaks are found, do not reuse or transport container, contact BAYER CROPSCIENCE LP at the number below for instructions. Disposing of Container: Do not reuse this container for any other purpose other than refilling (see preceding). Cleaning the container before final disposal is the responsibility of the person disposing of the container. To clean the container before final disposal, use the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration, and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour, or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling, if available, or puncture and dispose of in a sanitary landfill or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

Do not transport if container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire, or other emergency, contact BAYER CROPSCIENCE LP at 1-800-334-7577, day or night.

**CONDITIONS OF SALE AND LIMITATIONS OF WARRANTY
AND LIABILITY**

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, plant injury, other property damage, as well as other unintended consequences may result because of factors beyond the control of Bayer CropScience LP. Those factors include, but are not limited to, weather conditions, presence of other materials or the manner of use or application. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT.

For product information call: 1-800-331-2867

Bayer (reg'd), the Bayer Cross (reg'd) and Method® are registered trademarks of Bayer.

Produced for:
Bayer Environmental Science
A Division of Bayer CropScience LP
2 T. W. Alexander Drive
Research Triangle Park, NC 27709

Bayer

DO NOT USE PLANT MATERIAL TREATED WITH METHOD® 240SL HERBICIDE FOR MULCH OR COMPOST



Method® 240SL HERBICIDE

Soluble Liquid
For Non-Crop Use

ACTIVE INGREDIENT: By Weight

Potassium salt of aminocyclopyrachlor	
Potassium salt of 6-amino-5-chloro-2-cyclopropyl-4-pyrimidinencarboxylic acid*	25%
OTHER INGREDIENTS:	75%
TOTAL:	100%

*Acid Equivalent: 6-Amino-5-chloro-2-cyclopropyl-4-pyrimidinencarboxylic acid - 2 pounds acid per gallon or 21.2% JEPA Reg. No. 432-1565

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing.
PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers and loaders must wear:

Long-sleeved shirt and long pants. Shoes plus socks.

Applicators: After the product has been diluted in accordance with label directions for use, shirt, pants, socks, and shoes are sufficient Personal Protective Equipment (PPE).

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statement: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240 (d) (4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-334-7577 for emergency medical treatment information.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store product in original container only. Store in a cool, dry place.

Pesticide Disposal: Waste resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

Refer to the Net Contents section of this product's labeling for the applicable "Nonrefillable Container" or "Refillable Container" designation.

Nonrefillable Rigid Plastic and Metal Containers (Capacity Equal to or Less Than 5 Gallons):

Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. (continued)

continued

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Rigid Plastic and Metal Containers (Capacity Greater Than 5 Gallons): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Rigid Plastic and Metal Containers, e.g., Intermediate Bulk Containers (IBC) (Size or Shape Too Large to be Tipped, Rolled or Turned Upside Down): Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying the contents from this container into application equipment or mix tank and before final disposal using the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour, or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

All Refillable Containers: Refillable container. Refilling Container: Refill this container with METHOD® 240SL HERBICIDE containing aminocyclopyrachlor potassium salt only. Do not reuse this container for any other purpose. Cleaning before refilling is the responsibility of the refiller. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn out threads and closure devices. If damage is found, do not use container; contact BAYER CROPSCIENCE LP at the number below for instructions. Check for leaks after refilling and before transporting. If leaks are found, do not reuse or transport container, contact BAYER CROPSCIENCE LP at the number below for instructions. Disposing of Container: Do not reuse this container for any other purpose other than refilling (see preceding). Cleaning the container before final disposal is the responsibility of the person disposing of the container. To clean the container before final disposal, use the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration, and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour, or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling, if available, or puncture and dispose of in a sanitary landfill or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

Do not transport if container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire, or other emergency, contact BAYER CROPSCIENCE LP at 1-800-334-7577, day or night.

PULL HERE TO OPEN

Product of China

See attached leaflet for complete First Aid Instructions, Precautionary Statements, Directions for Use and Storage and Disposal Instructions.

Net Contents: **2.5 Gallons**

84099295 84942561A 150820AV1

Produced for:
Bayer Environmental Science
A Division of Bayer CropScience LP
2 T. W. Alexander Drive
Research Triangle Park, NC 27709

Bayer



0100785740387443