



TMDL SUMMARY

APPENDIX 6-11

Dyer River

WATERSHED DESCRIPTION

This **TMDL** applies to a 9.35 mile section of Dyer River, located in the Towns of Jefferson and Newcastle, Maine. The impaired segment of the Dyer River begins in the central portion of the watershed and flows south through a predominantly forested area, crossing Clary Road and County Road along the way. After crossing Jones Woods Road, the stream flows through agriculture and then into a wetland before crossing South Dyer Neck Road and joining the Sheepscot River. The Dyer River watershed covers an area of 27.63 square miles. The majority of the watershed is located within the Towns of Jefferson and Newcastle, however, smaller portions of the watershed lie within the surrounding towns of Whitefield and Somerville.

- Runoff from agricultural land located throughout the south central portion of the watershed are likely the largest source of **nonpoint source (NPS) pollution** to Dyer River. Runoff from cultivated lands, active hay lands, and grazing areas can transport nitrogen and phosphorus to the nearest section of the stream.
- The Dyer River watershed is predominately non-developed (97.5%). Forested areas (80.9%) within the watershed absorb and filter pollutants helping protect both water quality in the stream and stream channel stability. Wetlands (8.6%) may also help filter nutrients.
- Non-forested areas within the watershed are predominantly agricultural (7.33%) and are located in the south central portion of the watershed.
- Developed areas (2.5%) with impervious surfaces in close proximity to the stream may impact water quality.
- Dyer River is on Maine's 303(d) list of Impaired Streams (Maine DEP, 2013).

Definitions

- **Total Maximum Daily Load (TMDL)** represents the total amount of pollutants that a waterbody can receive and still meet water quality standards.
- **Nonpoint Source Pollution** refers to pollution that comes from many diffuse sources across the landscape, and is typically transported by rain or snowmelt runoff.

Waterbody Facts

Segment ID:

ME0105000305_528R03

Town: Jefferson and Newcastle, ME

County: Lincoln

Impaired Segment Length:
9.35 miles


Classification: Class B

Direct Watershed: 27.63 mi²
(17,683 acres)


Impairment Cause Listing:
Dissolved Oxygen

Watershed Agricultural Land Use: 7.33%

Major Drainage Basin:
Kennebec River



Watershed Land Uses






Figure 1: Land Use in the Dyer River Watershed

WHY IS A TMDL ASSESSMENT NEEDED

The Dyer River, a Class B freshwater stream, has been assessed by Maine DEP as not meeting water quality standards for the designated use of aquatic life, and placed on the 303(d) list of impaired waters under the Clean Water Act. The Clean Water Act requires that all 303(d)-listed waters undergo a TMDL assessment that describes the impairments and establishes a target to guide the measures needed to restore water quality. The goal is for all waterbodies to comply with state water quality standards.

Agriculture in the Dyer River watershed makes up about 7% of the total land area. This is almost three times the area of developed land, which accounts for 2.5% of land area. 31% of the impaired stream segment length passes through agricultural lands (Figure 1) therefore making agricultural land uses a likely contributor of sediment and nutrient enrichment to the stream. The close proximity of many agricultural lands to the stream further increases the likelihood that nutrients from disturbed soils, manure, and fertilizers will reach the stream. A large cattle operation located on Jones Woods Road and North Newcastle Road is known to spread liquid manure on all hay fields. Excess nutrient input into Dyer River through tributaries in the area is a great concern. However, the Dyer River watershed also has 8.6% wetlands, which may naturally lower dissolved oxygen concentrations in the portions of the river that flow through them.



*Dyer River downstream of the Jones Woods Road crossing. Station 917.
Photo: FB Environmental*

WATER QUALITY DATA ANALYSIS

Maine DEP uses a variety of data types to measure the ability of a stream to adequately support aquatic life, including; dissolved oxygen, benthic macroinvertebrates, and periphyton (algae). The aquatic life impairment in the Dyer River is based on historic dissolved oxygen data.

TMDL ASSESSMENT APPROACH: NUTRIENT MODELING OF IMPAIRED AND ATTAINMENT STREAMS

NPS pollution is difficult to measure directly, because it comes from many diffuse sources spread across the landscape. For this reason, a nutrient loading model, MapShed, was used to estimate the sources of pollution based on well-established hydrological equations; detailed maps of soil, land use, and slope; many years of daily weather data; and direct observations of agriculture and other land uses within the watershed.

The nutrient loading estimates for the impaired stream were compared to similar estimates for five non-impaired (attainment) streams of similar watershed land uses across the state. The TMDL for the impaired stream was set as the mean nutrient loading estimate of these attainment stream watersheds, and units of mass per unit watershed area per year (kg/ha/year) were used. The difference in loading estimates between the impaired and attainment watersheds represents the percent reduction in nutrient loading required under this TMDL. The attainment streams and their nutrient and sediment loading estimates and TMDL are presented below in Table 1.

Table 1: Numeric Targets for Pollutant Loading Based on MapShed Model Outputs for Attainment Streams

Attainment Streams	Town	TP load (kg/ha/yr)	TN load (kg/ha/yr)	Sediment load (1000 kg/ha/yr)
Martin Stream	Fairfield	0.14	3.4	0.008
Footman Brook	Exeter	0.33	6.4	0.058
Upper Kenduskeag Stream	Corinth	0.29	5.6	0.047
Upper Pleasant River	Gray	0.22	4.6	0.016
Moose Brook	Houlton	0.25	5.9	0.022
Total Maximum Daily Load		0.24	5.2	0.030

RAPID WATERSHED ASSESSMENT

Habitat Assessment

A Habitat Assessment survey was conducted on both the impaired and attainment stream. The assessment approach is based on the *Rapid Bioassessment Protocols for Use in Wadeable Streams and Rivers* (Barbour et al., 1999), which integrates various parameters relating to the structure of physical habitat. The habitat assessments include a general description of the site and physical characterization and visual assessment of in-stream and riparian habitat quality.

Based on Rapid Bioassessment protocols for low gradient streams, the Dyer River received a score of 155 out of a total 200 for quality of habitat. Higher scores indicate better habitat. The range of habitat assessment scores for attainment streams was 155 to 179.

Habitat assessments were conducted on a relatively short sample reach (about 100-200 meters for a typical small stream) near the most downstream Maine DEP sample station in the watershed. For both impaired and attainment streams, the assessment location was usually near a road crossing for ease of access. In the Dyer River watershed, the downstream sample station was located in a forested portion of the stream with a thick buffer in most areas. A maintained lawn is located adjacent to the stream with a minimal buffer on the east side of the sample reach on the north side of Jones Woods Road.

Figure 2 (right) shows the range of habitat assessment scores for all attainment and impaired streams, as well as for the Dyer River. The overlapping attainment and impaired stream scores indicate that factors other than habitat should be considered when addressing the impairments in the Dyer River. Consideration should be given to major “hot spots” in the Dyer River watershed as potential sources of NPS pollution contributing to the water quality impairment.




Figure 2: Habitat Assessment Scores

Pollution Source Identification

Pollution source identification assessments were conducted for both the Dyer River (impaired) and the attainment streams. The source identification work is based on an abbreviated version of the Center for Watershed Protection’s Unified Subwatershed and Site Reconnaissance method (Wright, et al., 2005). The abbreviated method includes both a desktop and field component. The desktop assessment consists of generating and reviewing maps of the watershed boundary, roads, land use and satellite imagery, and then identifying potential NPS pollution locations, such as road crossings, agricultural fields, and large areas of bare soil. When available, multiple sources of satellite imagery were reviewed. Occasionally,

the high resolution of the imagery allowed for observations of livestock, row crops, eroding stream banks, sediment laden water, junkyards, and other potential NPS concerns that could affect stream quality. As many potential pollution sources as possible were visited, assessed and documented in the field. Field visits were limited to NPS sites that were visible from roads or a short walk from a roadway. Neighborhoods were assessed for NPS pollution at the whole neighborhood level including streets and storm drains (where applicable). The assessment does not include a scoring component, but does include a detailed summary of findings and a map indicating documented NPS sites throughout the watershed.

The watershed source assessment for the Dyer River was completed on July 3, 2012. In-field observations of erosion, lack of vegetated stream buffer, extensive impervious surfaces, high-density neighborhoods and agricultural activities were documented throughout the watershed (Table 2, Figure 3).

Table 2: Pollution Source ID Assessment for the Dyer River Watershed

Potential Source			Notes
ID#	Location	Type	
1	North Dyer Neck Road	Agriculture	<ul style="list-style-type: none"> Active pasture. Manure smell.
2	Jones Woods Road	Road Crossing	<ul style="list-style-type: none"> Sample reach location. DEP Biomonitoring Station 917 upstream of crossing. Mowed lawns adjacent to stream.
3	Jones Woods Road/North Newcastle Road	Agriculture	<ul style="list-style-type: none"> Large dairy farm with over 300 cows. Large open graze land. Strong manure odor throughout this area. Impounded tributaries near pasture with algal growth. Tributaries flowing through graze land; potential direct access to stream by cows.
6	Between South Dyer Neck Road & North Newcastle Road	Wetland	<ul style="list-style-type: none"> Large wetland complex. Obvious source of low dissolved oxygen.
15	Atkins Road/South Clary Road	Gravel pit	<ul style="list-style-type: none"> Gravel Products, LLC Pit #3; large pit. Seems active.
16	South Clary Road	Agriculture	<ul style="list-style-type: none"> Active row crops and hay fields.
18	South Clary Road and Banks Lane	Agriculture	<ul style="list-style-type: none"> Small farm: row crops and hay fields.
20	South Clary Road	Gravel Pit	<ul style="list-style-type: none"> Sherwood Wood Products Gravel Pit; large pit. Seems active.
21	Road Crossing	South Clary Road	<ul style="list-style-type: none"> Point bars and deposits visible from roadway. Braided channel in some parts.




Figure 3b: Source ID Locations in the Upper and Lower Dyer River Watershed. Note that this map only shows the lower portion of the impaired stream and watershed.

NUTRIENT LOADING – MAPSHED ANALYSIS

The MapShed model was used to estimate stream loading of sediment, total nitrogen and total phosphorus in Dyer River (impaired) plus five attainment watersheds throughout the state.. The model estimated nutrient loads over a 15-year period (1990-2004), which was determined by the available weather data provided within MapShed. This extended period captures a wide range of hydrologic conditions to account for variations in nutrient and sediment loading over time.

Many quality assured and regionally calibrated input parameters are provided with MapShed. Additional input parameters were manually entered into the model based on desktop research and field observations, as described in the sections on Habitat Assessment and Pollution Source Identification. These manually adjusted parameters included estimates of livestock animal units, agricultural stream miles with intact vegetative buffer, Best Management Practices (BMPs), and estimated wetland retention and/or drainage areas.

Livestock Estimates

Livestock waste contains nutrients which can cause water quality impairment. The nutrient loading model considers numbers and types of animals. Table 3 (right) provides estimates of livestock (numbers of animals) in the watershed, based on direct observations made in the watershed, plus other publicly available data.

The Dyer River watershed is predominantly forested, with substantial mixed agricultural land uses as well. In the upper watershed, small crops were grown in hobby farm operations. Hay fields were most common. In the lower watershed pasture and hay fields dominate. A very large dairy farm is located on Jones Woods Road. An estimated 300 cows are located here. Large pastures were observed with grazing cows and a strong manure smell was documented. Three tributaries drain these fields into the Dyer River, and multiple impoundments on these tributaries were green with significant algal growth. A conversation with a neighbor during our habitat assessment indicated the use of liquid manure on the dairy’s fields. This farm is considered a hot spot and a clear source of NPS pollution to the Dyer River. About 10 alpacas were also observed along North Newcastle Road. This hobby farm was not in close proximity to the Dyer River.

Table 3: Livestock Estimates in the Dyer River Watershed

Type	Dyer River
Dairy Cows	300
Beef Cows	
Broilers	
Layers	
Hogs/Swine	
Sheep	
Horses	
Turkeys	
Other	10 (alpacas)
Total	310

Vegetated Stream Buffer in Agricultural Areas

Vegetated stream buffers are areas of trees, shrubs, and/or grasses adjacent to streams, lakes, ponds or wetlands which provide nutrient loading attenuation (Evans & Corradini, 2012). MapShed considers natural vegetated stream buffers within agricultural areas as providing nutrient load attenuation. The width of buffer strips is not defined within the MapShed manual, and was considered to be 75 feet for this analysis. Geographic Information System (GIS) analysis of recent aerial photos along with field reconnaissance observations were used to estimate the number of agricultural stream miles with and without vegetative buffers, and these estimates were directly entered into the model.

The Dyer River is a 9.4 mile-long impaired segment as listed by Maine DEP. As modeled, the total stream miles (including tributaries) within the watershed was calculated as 74.9 miles. Of this total, 2.9 stream miles are located within agricultural areas; of this length, 1.4 miles (48%) show a 75-foot or greater vegetated buffer (Table 4, Fig. 4). By contrast, agricultural stream miles (as modeled) with a 75-foot vegetated buffer in the attainment stream watersheds ranged from 34% to 92%, with an average of 61%.

Table 4: Summary of Vegetated Buffers in Agricultural Areas

Dyer River
<ul style="list-style-type: none"> • 74.9 stream miles in watershed (includes ephemeral streams) • 2.9 stream miles in agricultural areas • 48% of agricultural stream miles have a vegetated buffer




Figure 4: Agricultural Stream Buffer in the Dyer River Watershed

Best Management Practices (BMPs)

For this modeling effort, four commonly used BMPs were entered based on literature values. These estimates were applied equally to impaired and attainment stream watersheds. More localized data on agricultural practices would improve this component of the model.

- *Cover Crops*: Cover crops are the use of annual or perennial crops to protect soil from erosion during time periods between harvesting and planting of the primary crop. The percent of agricultural acres cover crops used within the model is estimated at 4%. This figure is based on information from the 2007 USDA Census stating that 4.1% of cropland acres is left idle or used for cover crops or soil improvement activity, and not pastured or grazed (USDA, 2007b).
- *Conservation Tillage*: Conservation tillage is any kind of system that leaves at least 30% of the soil surface covered with crop residue after planting. This reduces soil erosion and runoff and is one of the most commonly used BMPs. This BMP was assumed to occur in 42% of agricultural land. This figure is based on a number given by the Conservation Tillage Information Center's 2008 Crop Residue Management Survey stating that 41.5% of U.S. acres are currently in conservation tillage (CTIC, 2000).
- *Strip Cropping / Contour Farming*: This BMP involves tilling, planting and harvesting perpendicular to the gradient of a hill or slope using high levels of plant residue to reduce soil erosion from runoff. This BMP was assumed to occur in 38% of agricultural lands, based on a study done at the University of Maryland (Lichtenberg, 1996).
- *Grazing Land Management*: This BMP consists of ensuring adequate vegetation cover on grazed lands to prevent soil erosion from overgrazing or other forms of over-use. This usually employs a rotational grazing system where hays or legumes are planted for feed and livestock is rotated through several fenced pastures. In this TMDL, a figure of 75% of hay and pasture land is assumed to utilize grazing land management. This figure is based on a study by Farm Environmental Management Systems of farming operations in Canada (Rothwell, 2005).

Pollutant Load Attenuation by Lakes, Ponds and Wetlands

Depositional environments such as ponds and wetlands can attenuate watershed sediment loading. This information is entered into the nutrient loading model by a simple percentage of watershed area draining to a pond or a wetland. The Dyer River watershed is 9% wetland, and overall 30% of the watershed drains to wetlands. Percent of watershed draining to a wetland in the attainment watersheds ranged from 15% to 60%, with an average of 35%.

NUTRIENT MODELING RESULTS

The MapShed model simulates surface runoff using daily weather inputs of rainfall and temperature. Erosion and sediment yields are estimated using monthly erosion calculations and land use/soil composition values for each source area. Below, selected results from the watershed loading model are presented. The TMDL itself is expressed in units of kilograms per hectare per year. The additional results shown below assist in better understanding the likely sources of pollution. The model results for the Dyer River indicate that no reductions of sediment or nutrients are needed to improve water quality. Below, loading for sediment, nitrogen and phosphorus are discussed individually.

Sediment

Sediment loading in the Dyer River watershed is primarily attributed to crop land which accounts for 40% of the total sediment load. Agricultural sources combined (including crop land) attribute over half of the sediment loading in the Dyer River watershed (Table 5 and Figure 5). High density development and forested land also contribute significantly to the sediment load at 22% and 21% of the total load, respectively. Note that total loads by mass cannot be directly compared between watersheds due to differences in watershed area. See section *TMDL: Target Nutrient Levels for the Dyer River* below for loading estimates that have been normalized by watershed area.

Table 5: Total Sediment Loads by Source

Dyer River	Sediment (1000kg/year)	Sediment (%)
Source Load		
<i>Hay/Pasture</i>	7.97	14%
<i>Crop land</i>	23.57	40%
<i>Forest</i>	12.12	21%
<i>Wetland</i>	0.40	1%
<i>Disturbed Land</i>	0	0%
<i>Sandy Areas</i>	0.08	0%
<i>Low Density Mixed</i>	0.78	1%
<i>Medium Density Mixed</i>	0	0%
<i>High Density Mixed</i>	12.85	22%
<i>Low Density Residential</i>	0.71	1%
<i>Medium Density Residential</i>	0	0%
<i>High Density Residential</i>	0	0%
<i>Farm Animals</i>	0	0%
<i>Septic Systems</i>	0	0%
Source Load Total:	58.48	100%
Pathway Load		
<i>Stream Banks</i>	45.10	-
<i>Subsurface / Groundwater</i>	0	-
Total Watershed Mass Load:		
	103.58	




Figure 5: Total Sediment Loads by Source in the Dyer River Watershed

Total Nitrogen

Nitrogen loading in the Dyer River watershed is attributed primarily to farm animals, making up 42% of the total load. Agricultural sources combined (including farm animals) account for almost 60% of the nitrogen load in the Dyer River watershed. Forested land contributes 21% of the total load. Table 6 and Figure 6 show the estimated total nitrogen load in terms of mass and percent of total and by source. Note that total loads by mass cannot be directly compared between watersheds due to differences in watershed area. See section *TMDL: Target Nutrient Levels for the Dyer River* below for loading estimates that have been normalized by watershed area.

Table 6: Total Nitrogen Loads by Source

Dyer River	Total N (kg/year)	Total N (%)
Source Load		
Hay/Pasture	359.9	5%
Crop land	727.9	11%
Forest	1377.7	21%
Wetland	364.8	5%
Disturbed Land	0	0%
Sandy Areas	0.2	0%
Low Density Mixed	24.3	0%
Medium Density Mixed	0	0%
High Density Mixed	643.3	10%
Low Density Residential	21.9	0%
Medium Density Residential	0	0%
High Density Residential	0	0%
Farm Animals	2795.6	42%
Septic Systems	345.5	5%
Source Load Total:	6661.2	100%
Pathway Load		
Stream Banks	37.6	-
Subsurface / Groundwater	23019.5	-
Total Watershed Mass Load:	29718.3	




Figure 6: Total Nitrogen Loads by Source in the Dyer River Watershed

Total Phosphorus

Phosphorus loading in the Dyer River watershed is attributed primarily to farm animals, making up almost 60% of the total phosphorus load. Agricultural sources combined (including farm animals) account for 80% of the total load to the Dyer River. Phosphorus loads are presented in Table 7 and Figure 7. Note that total loads by mass cannot be directly compared between watersheds due to differences in watershed area. See section *TMDL: Target Nutrient Levels for the Dyer River* below for loading estimates that have been normalized by watershed area

Table 7: Total Phosphorus Loads by Source

Dyer River	Total P (kg/year)	Total P (%)
Source Load		
<i>Hay/Pasture</i>	116.7	13%
<i>Crop land</i>	71.7	8%
<i>Forest</i>	76.5	9%
<i>Wetland</i>	18.4	2%
<i>Disturbed Land</i>	0	0%
<i>Sandy Areas</i>	0.1	0%
<i>Low Density Mixed</i>	2.6	0%
<i>Medium Density Mixed</i>	0	0%
<i>High Density Mixed</i>	63.9	7%
<i>Low Density Residential</i>	2.3	0%
<i>Medium Density Residential</i>	0	0%
<i>High Density Residential</i>	0	0%
<i>Farm Animals</i>	518.7	59%
<i>Septic Systems</i>	3.1	0%
Source Load Total:	874.0	100%
Pathway Load		
<i>Stream Banks</i>	11.0	-
<i>Subsurface / Groundwater</i>	781.0	-
Total Watershed Mass Load:	1665.9	




Figure 7: Total Phosphorus Loads by Source in the Dyer River Watershed

TMDL: TARGET NUTRIENT LEVELS FOR THE DYER RIVER

The existing loads for sediments and nutrients in the impaired segment of the Dyer River are listed in Table 8, along with the TMDL numeric target which was calculated from the average loading estimates of five attainment watersheds throughout the state. Table 9 presents a more detailed view of the modeling results and calculations used in Table 8 to define TMDL reductions, and compares the existing sediment and nutrient loads in the Dyer River to TMDL endpoints derived from the attainment waterbodies. An annual time frame provides a mechanism to address the daily and seasonal variability associated with nonpoint source loads.

Table 8: TMDL Targets Compared to Dyer River Pollutant Loading

TMDL POLLUTANT LOADS Annual Loads per Unit Area	Estimated Loads Dyer River	Total Maximum Daily Load Numeric Target	TMDL % REDUCTIONS Dyer River
<i>Sediment Load (1000 kg/ha/year)</i>	0.015	0.030	No Reduction Needed
<i>Nitrogen Load (kg/ha/year)</i>	4.31	5.2	No Reduction Needed
<i>Phosphorus Load (kg/ha/year)</i>	0.24	0.24	No Reduction Needed

Future Loading

The prescribed reduction in pollutants discussed in this TMDL reflects reduction from estimated existing conditions. Expansion of agricultural and development activities have the potential to increase runoff and associated pollutant loads to the Dyer River. To ensure that the TMDL targets are attained, future agriculture or development activities in the watershed will need to meet the TMDL targets. Future growth from population increases is a moderate threat in the Dyer River watershed because Lincoln County has increasing population trends, with a 3% increase between 2000 and 2008 (USM MSAC, 2009). The growth in agricultural lands is also increasing, with a 24% increase in the total number of farms in Lincoln County between 2002 and 2007. However, a decrease of 2% was seen in the land (acres) in farms between 2002 and 2007, and a 21% decrease occurred in the average farm size in this time period as well (USDA, 2007a). Future activities and BMPs that achieve TMDL reductions are addressed below.

Next Steps

The use of agricultural and developed area BMPs can reduce sources of polluted runoff in the Dyer River. It is recommended that municipal officials, landowners, and conservation stakeholders in Jefferson and Newcastle work together to develop a watershed management plan to:

- Encourage greater citizen involvement through the development of a watershed coalition to ensure the long term protection of the Dyer River;
- Address existing nonpoint source problems in the Dyer River watershed by instituting BMPs where necessary; and

- Prevent future degradation of the Dyer River through the development and/or strengthening of a local Nutrient Management Ordinance.

The Dyer River watershed is located within the larger Sheepscot River watershed which currently has two main associations focused on improving water quality within the greater watershed. The Sheepscot Valley Conservation Association and the Sheepscot River Watershed Coalition teamed up in 2011 and acquired an EPA 319 grant used to survey the Dyer River watershed. Bacteria sampling and an NPS survey were conducted in the summer of 2011. Recommendations following the survey included monitoring river setbacks from forestry and agricultural land uses, maintaining a long-term bacteria sampling program, investigating additional non-agricultural bacteria sources, providing outreach to state agencies that monitor gravel pits, and engaging the local community in conservation and awareness of environmental issues in the Dyer River watershed (Baeder, 2011).

Table 9: Modeling Results Calculations for Derived Numeric Targets and Reduction Loads for Dyer River

Dyer River				
	Area ha	Sediment 1000kg/yr	TN kg/yr	TP kg/yr
Land Uses				
<i>Hay/Pasture</i>	399	8.0	360.0	116.7
<i>Crop land</i>	131	23.6	728.0	71.7
<i>Forest</i>	5556	12.1	1377.7	76.5
<i>Wetland</i>	591	0.4	364.8	18.4
<i>Disturbed Land</i>	0	0.0	0.0	0.0
<i>Sandy Areas</i>	28	0.1	0.2	0.1
<i>Low Density Mixed</i>	30	0.8	24.3	2.6
<i>High Density Mixed</i>	139	12.9	643.3	63.9
<i>Low Density Residential</i>	27	0.7	21.9	2.3
Other Sources				
<i>Farm Animals</i>			2795.6	518.7
<i>Septic Systems</i>			345.5	3.07
Pathway Loads				
<i>Stream Banks</i>		45.1	37.6	11.0
<i>Groundwater</i>			23019.5	781.0
Total Annual Load		104 x 1000 kg	29718 kg	1666.0 kg
Total Area	6901 ha			
Total Maximum Daily Load		0.015 1000kg/ha/year	4.31 kg/ha/year	0.24 kg/ha/year

REFERENCES

- Barbour, M.T., J. Gerritsen, B.D. Snyder, and J.B. Stribling. 1999. Rapid Bioassessment Protocols for Use in Wadeable Streams and Rivers: Periphyton, Benthic Macroinvertebrates and Fish, Second Edition. EPA 841-B-99-002. U.S. Environmental Protection Agency; Office of Water; Washington, D.C.
- Baeder, C. & M. Hoffman. 2011. Dyer River Watershed Bacteria and NPS Survey Report. Sheepscot Valley Conservation Association. Newcastle, Maine.
- Conservation Tillage Information Center (CTIC). 2000. Crop Residue Management Survey. National Association of Conservation Districts. Retrieved from: <http://www.ctic.purdue.edu/CTIC/CTIC.html>.
- Davies, S. P., and L. Tsomides. 2002. Methods for Biological Sampling of Maine's Rivers and Streams. DEP LW0387-B2002, Maine Department of Environmental Protection, Augusta, ME.
- Evans, B.M., & Corradini, K.J. 2012. MapShed Version 1.0 Users Guide. Penn State Institute of Energy and the Environment. Retrieved from: <http://www.ctic.purdue.edu>.
- Lichtenberg, E. 1996. Using Soil and Water Conservation Practices to Reduce Bay Nutrients: How has Agriculture Done? Economic Viewpoints. Maryland Cooperative Extension Service, University of Maryland at College Park and University of Maryland Eastern Shore, Department of Agricultural and Resource Economics, 1(2).
- Maine Department of Environmental Protection (Maine DEP). 2013. Draft 2012 Integrated Water Quality Monitoring and Assessment Report. Bureau of Land and Water Quality, Augusta, ME.
- Rothwell, N. 2005. Grazing Management in Canada. Farm Environmental Management in Canada. <http://publications.gc.ca/Collection/Statcan/21-021-M/21-021-MIE2005001.pdf>.
- University of Southern Maine Muskie School of Public Service, Maine Statistical Analysis Center (USM MSAC). December, 2009. Retrieved from: <http://muskie.usm.maine.edu/justiceresearch/Publications/County/Lincoln.pdf>
- United States Department of Agriculture (USDA). 2007a. 2007 Census of Agriculture: Lincoln County, Maine. Retrieved from: http://www.agcensus.usda.gov/Publications/2007/Online_Highlights/County_Profiles/Maine/cp23015.pdf
- United States Department of Agriculture (USDA). 2007b. 2007 Census of Agriculture: State and County Reports. National Agricultural Statistics Service. Retrieved from: http://www.agcensus.usda.gov/Publications/2007/Full_Report/Volume_1,_Chapter_1_State_Level/Maine/st23_1_008_008.pdf
- Wright, T., C. Swann, K. Cappiella, and T. Schueler. 2005. Unified Subwatershed and Site Reconnaissance: A User's Manual. Center for Watershed Protection. Ellicott City, MD.