

Antigen (Ag) testing is less sensitive than PCR. Therefore, if utilizing antigen testing, the recommendation is to serial antigen test at no less than twice a week, typically 3-4 days apart.

Example:

		PCR only	Antigen Only	Mix of PCR and Antigen
Once a Week	Week 1	Monday (PCR)	Monday (Ag) & Thursday (Ag)	Monday (PCR)
	Week 2	Monday (PCR)	Monday (Ag) & Thursday (Ag)	Monday (Ag) & Thursday (Ag)
	Week 3	Monday (PCR)	Monday (Ag) & Thursday (Ag)	Monday (PCR)
	Week 4	Monday (PCR)	Monday (Ag) & Thursday (Ag)	Monday (Ag) & Thursday (Ag)
Twice a week	Week 1	Monday (PCR) and Thursday (PCR)	Monday (Ag) & Thursday (Ag)	Monday (PCR) and Thursday (PCR)
	Week 2	Monday (PCR) and Thursday (PCR)	Monday (Ag) & Thursday (Ag)	Monday (Ag) & Thursday (Ag)
	Week 3	Monday (PCR) and Thursday (PCR)	Monday (Ag) & Thursday (Ag)	Monday (PCR) and Thursday (PCR)
	Week 4	Monday (PCR) and Thursday (PCR)	Monday (Ag) & Thursday (Ag)	Monday (Ag) & Thursday (Ag)

Last updated: 09/17/2021