

Maine Shared Community Health Needs Assessment: 2015

U.S. Data Sources

Indicator	Data Source	Year(s)	Other Notes
Demographics			
Population	U.S. Census	2014	2013 data was used for all age, racial and ethnic groups.
Population with a disability	U.S. Census	2010	Adults reporting any one of the six disability types are considered to have a disability: hearing difficulty, vision difficulty, cognitive difficulty, ambulatory difficulty, self-care difficulty, independent living difficulty.
Population density	U.S. Census	2010	Based on 2010 U.S. Census population.
Socioeconomic Status Measures			
Individuals living in poverty	U.S. Census	2009-2013	The poverty status of the household is determined by the poverty status of the householder. Households are classified as poor when the total income of the householder's family is below the appropriate poverty threshold. The American Community Survey measures poverty in the previous 12 months instead of the previous calendar year.
Children living in poverty	U.S. Census	2009-2013	The poverty status of the household is determined by the poverty status of the householder. Households are classified as poor when the total income of the householder's family is below the appropriate poverty threshold. The American Community Survey measures poverty in the previous 12 months instead of the previous calendar year.
High school graduation rate	U.S. Department of Education, National Center for Education Statistics	2012-13 School Year	Proportion of students who graduate with a regular diploma four years after starting ninth grade. Graduation rates include all public schools and all private schools that have 60% or more publicly funded students.
Median household income	U.S. Census	2009-2013	In 2013 inflation-adjusted dollars. This includes the income of the householder and all other individuals 15 years old and older in the household, whether they are related to the householder or not.
Percentage of people living in rural areas	NA	NA	Data not available.
Single-parent families	U.S. Census	2013	Families consist of a householder and one or more other people related to the householder by birth, marriage, or adoption. They do not include same-sex married couples even if the marriage was performed in a state issuing marriage certificates for same-sex couples. "Householder without a spouse present" is defined as a male householder without a wife present or a female householder without a husband present.
Unemployment rate	Bureau of Labor Statistics	2014	Unemployment rate of the civilian noninstitutionalized population averaged for the full year of 2014.
65+ living alone	U.S. Census	2013	Estimated number of one-person households with a person 65 years and older.
General Health Status			
Adults who rate their health fair to poor	BRFSS	2013	Adults rating their health as fair or poor vs. excellent, very good or good.
Adults with 14+ days lost due to poor mental health	NA	NA	Data not available.
Adults with 14+ days lost due to poor physical health	NA	NA	Data not available.
Adults with three or more chronic conditions	NA	NA	Data not available.

Maine Shared Community Health Needs Assessment: 2015

U.S. Data Sources

Indicator	Data Source	Year(s)	Other Notes
Mortality			
Life expectancy (Female)	CDC/National Center for Health Statistics	2012	Life expectancy at birth.
Life expectancy (Male)	CDC/National Center for Health Statistics	2012	Life expectancy at birth.
Overall mortality rate per 100,000 population	CDC/National Center for Health Statistics, National Vital Statistics System	2013	All deaths are defined as deaths in which the underlying cause of death was coded as ICD-10 any listed.
Access			
Adults with a usual primary care provider	BRFSS	2013	Adults that have one or more person they think of as their personal doctor or health care provider.
Individuals who are unable to obtain or delay obtaining necessary medical care due to cost	BRFSS	2013	Adults reporting that there was a time during the last 12 months when they needed to see a doctor but could not because of the cost.
Medicaid enrollment	U.S. Department of Health & Human Services, Centers for Medicare & Medicaid Services	2013	The percent of individuals participating in Medicaid.
Percent of children ages 0-19 enrolled in Medicaid	U.S. Department of Health & Human Services, Centers for Medicare & Medicaid Services	2013	The percent of individuals participating in Medicaid.
Percent uninsured	U.S. Census	2014	Estimated number of U.S. people who do not currently have health insurance.
Health Care Quality			
Ambulatory care-sensitive condition hospital admission rate per 100,000 population	Agency for Healthcare Research and Quality	2012	PQI = Prevention Quality Indicators, a set of measures that can be used with hospital inpatient discharge data to identify quality of care for ambulatory care-sensitive conditions. Additional information at: AHRQ Quality Indicators, Version 4.4, Agency for Healthcare Research and Quality: U.S. Department of Health and Human Services. http://www.qualityindicators.ahrq.gov .
Ambulatory care-sensitive condition emergency department rate per 100,000 population	NA	NA	Data not available.
Oral Health			
Adults with visits to a dentist in the past 12 months	BRFSS	2012	Adults who last visited the dentist or a dental clinic for any reason in the past 12 months.
Medicaid members under 18 with a visit to the dentist in the past year	NA	NA	Data not available.
Respiratory			
Asthma emergency department visits per 10,000 population	NA	NA	Data not available.
COPD diagnosed	BRFSS	2013	Adults that have been told by a doctor, nurse or health professional that they have COPD chronic obstructive pulmonary disease, emphysema, or chronic bronchitis.
COPD hospitalizations per 100,000 population	NA	NA	Data not available.
Current asthma (Adults)	BRFSS	2013	Adults that have been told by a doctor, nurse or health professional that they had asthma and that they still have asthma.
Current asthma (Youth 0-17)	BRFSS	2013	Children that have been told by a doctor, nurse or health professional that they had asthma and that they still have asthma.
Pneumonia emergency department rate per 100,000 population	NA	NA	Data not available.

Maine Shared Community Health Needs Assessment: 2015

U.S. Data Sources

Indicator	Data Source	Year(s)	Other Notes
Pneumonia hospitalizations per 100,000 population	NA	NA	Data not available.
Cancer			
Mortality – all cancers per 100,000 population	SEER	2011	All cancer: SEER Cause of Death Recode: 20010-37000 (which include ICD-10 codes: C00-C97).
Incidence – all cancers per 100,000 population	SEER	2009-2011	All cancer: SEER Site Recode: 20010-37000 (which include ICD-O-3 codes: C00-C797).
Bladder cancer incidence per 100,000 population	SEER	2009-2011	Cancer Incidence: The number of people who develop cancer (new cancer cases) during a specified period of time in a specified population. Incidence case definitions exclude histologies consistent with Kaposi sarcoma and mesothelioma, where applicable.
Female breast cancer mortality per 100,000 population	SEER	2011	Cancer Deaths: Deaths with malignant cancer as the underlying cause of death.
Breast cancer late-stage incidence (females only) per 100,000 population	SEER	2009-2011	Cancer Incidence: The number of people who develop cancer (new cancer cases) during a specified period of time in a specified population. Incidence case definitions exclude histologies consistent with Kaposi sarcoma and mesothelioma, where applicable.
Female breast cancer incidence per 100,000 population	SEER	2009-2011	Cancer Incidence: The number of people who develop cancer (new cancer cases) during a specified period of time in a specified population. Incidence case definitions exclude histologies consistent with Kaposi sarcoma and mesothelioma, where applicable.
Mammograms females age 50+ in past two years	BRFSS	2012	Females ages 50 years and older who reported they had a mammogram within the past 2 years.
Colorectal cancer mortality per 100,000 population	SEER	2011	Cancer Deaths: Deaths with malignant cancer as the underlying cause of death.
Colorectal late-stage incidence per 100,000 population	SEER	2009-2011	Cancer Incidence: The number of people who develop cancer (new cancer cases) during a specified period of time in a specified population. Incidence case definitions exclude histologies consistent with Kaposi sarcoma and mesothelioma, where applicable.
Colorectal cancer incidence per 100,000 population	SEER	2009-2011	Cancer Incidence: The number of people who develop cancer (new cancer cases) during a specified period of time in a specified population. Incidence case definitions exclude histologies consistent with Kaposi sarcoma and mesothelioma, where applicable.
Colorectal screening	BRFSS	2013	Adults ages 50 years and older who reported that they had a home blood stool test (e.g., FOBT or FIT) within the past year OR sigmoidoscopy within the past 5 years and home blood stool test within the past 3 years OR a colonoscopy within the past 10 years.
Lung cancer mortality per 100,000 population	SEER	2011	Cancer Deaths: Deaths with malignant cancer as the underlying cause of death.
Lung cancer incidence per 100,000 population	SEER	2009-2011	Cancer Incidence: The number of people who develop cancer (new cancer cases) during a specified period of time in a specified population. Incidence case definitions exclude histologies consistent with Kaposi sarcoma and mesothelioma, where applicable.

Maine Shared Community Health Needs Assessment: 2015

U.S. Data Sources

Indicator	Data Source	Year(s)	Other Notes
Melanoma incidence per 100,000 population	SEER	2011	Cancer Incidence: The number of people who develop cancer (new cancer cases) during a specified period of time in a specified population. Incidence case definitions exclude histologies consistent with Kaposi sarcoma and mesothelioma, where applicable.
Pap smears females ages 21-65 in past three years	BRFSS	2012	Females with intact cervix, that have received a pap smear within the past three years.
Prostate cancer mortality per 100,000 population	SEER	2011	Cancer Deaths: Deaths with malignant cancer as the underlying cause of death.
Prostate cancer incidence per 100,000 population	SEER	2009-2011	Cancer Incidence: The number of people who develop cancer (new cancer cases) during a specified period of time in a specified population. Incidence case definitions exclude histologies consistent with Kaposi sarcoma and mesothelioma, where applicable.
Tobacco-related neoplasms, mortality per 100,000 population	SEER	2011	Cancer Deaths: Deaths with malignant cancer as the underlying cause of death.
Tobacco-related neoplasms, incidence per 100,000 population	SEER	2009-2011	Cancer Incidence: The number of people who develop cancer (new cancer cases) during a specified period of time in a specified population. Incidence case definitions exclude histologies consistent with Kaposi sarcoma and mesothelioma, where applicable.
Cardiovascular Disease			
Acute myocardial infarction hospitalizations per 10,000 population	NA	NA	Data not available.
Acute myocardial infarction mortality per 100,000 population	CDC/National Center for Health Statistics, National Vital Statistics System	2013	ICD-10 I21-I22
Cholesterol checked every five years	BRFSS	2013	Adults reporting that they last had their blood cholesterol checked within the past 5 years.
Coronary heart disease mortality per 100,000 population	CDC/National Center for Health Statistics, National Vital Statistics System	2013	ICD-10 I20-I25
Heart failure hospitalizations per 10,000 population	NA	NA	Data not available.
Hypertension prevalence	BRFSS	2013	Adults who have ever been told by a doctor, nurse, or other health professional that they have high blood pressure.
High cholesterol	BRFSS	2013	Adults who have been told by a doctor or other health professional that their blood cholesterol is high.
Hypertension hospitalizations per 100,000 population	NA	NA	Data not available.
Stroke hospitalizations per 10,000 population	NA	NA	Data not available.
Stroke mortality per 100,000 population	CDC/National Center for Health Statistics, National Vital Statistics System	2013	ICD-10 I60-I69
Diabetes			
Diabetes prevalence (ever been told)	BRFSS	2013	Adults that have ever been told by a doctor or other health professional that they have diabetes.
Pre-diabetes prevalence	NA	NA	Data not available.
Adults with diabetes who have eye exam annually	NA	NA	Data not available.
Adults with diabetes who have foot exam annually	NA	NA	Data not available.

Maine Shared Community Health Needs Assessment: 2015

U.S. Data Sources

Indicator	Data Source	Year(s)	Other Notes
Adults with diabetes who have had an A1C test twice per year	NA	NA	Data not available.
Adults with diabetes who have received formal diabetes education	BRFSS	2013	Adults with diabetes who have ever taken a course or class in how to manage your diabetes themselves.
Diabetes emergency department visits (principal diagnosis) per 100,000 population	NA	NA	Data not available.
Diabetes hospitalizations (principal diagnosis) per 10,000 population	NA	NA	Data not available.
Diabetes long-term complication hospitalizations	NA	NA	Data not available.
Diabetes mortality (underlying cause) per 100,000 population	CDC/National Center for Health Statistics, National Vital Statistics System	2013	ICD-10 E10-E14
Environmental Health			
Children with confirmed elevated blood lead levels (% among those screened)	NA	NA	Data not available.
Children with unconfirmed elevated blood lead levels (% among those screened)	NA	NA	Data not available.
Homes with private wells tested for arsenic	NA	NA	Data not available.
Lead screening among children age 12-23 months	NA	NA	Data not available.
Lead screening among children age 24-35 months	NA	NA	Data not available.
Immunization			
Adults immunized annually for influenza	NA	NA	Data not available.
Adults immunized for pneumococcal pneumonia (ages 65 and older)	BRFSS	2013	Risk factor for adults aged 65 or older that have ever had a pneumonia shot.
Immunization exemptions among kindergarteners for philosophical reasons	NA	NA	Data not available.
Two-year-olds up to date with "Series of Seven Immunizations" 4-3-1-3-3-1-4	NA	NA	Data not available.
Infectious Disease			
Hepatitis A (acute) incidence per 100,000 population	CDC/Division of Viral Hepatitis and National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention	2014	Defined as the number of new infections during 2014.
Hepatitis B (acute) incidence per 100,000 population	CDC/Division of Viral Hepatitis and National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention	2014	Defined as the number of new infections during 2014.
Hepatitis C (acute) incidence per 100,000 population	CDC/Division of Viral Hepatitis and National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention	2014	Defined as the number of new infections during 2014.
Incidence of past or present hepatitis C virus (HCV) per 100,000 population	NA	NA	Data not available.
Incidence of newly reported chronic hepatitis B virus (HBV) per 100,000 population	NA	NA	Data not available.

Maine Shared Community Health Needs Assessment: 2015

U.S. Data Sources

Indicator	Data Source	Year(s)	Other Notes
Lyme disease incidence per 100,000 population	Centers for Disease Control and Prevention	2014	Defined as the number of new infections during 2014.
Pertussis incidence per 100,000 population	Centers for Disease Control and Prevention	2014	Incidence is defined as the number of new infections during 2014.
Tuberculosis incidence per 100,000 population	Centers for Disease Control and Prevention	2014	New diagnoses, regardless of when infection occurred or stage of disease at diagnosis.
STD/HIV			
AIDS incidence per 100,000 population	CDC/Division of STD Prevention, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention	2014	Incidence is defined as the number of new infections during 2014.
Chlamydia incidence per 100,000 population	CDC/Division of STD Prevention, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention	2014	Incidence is defined as the number of new infections during 2014.
Gonorrhea incidence per 100,000 population	CDC/Division of STD Prevention, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention	2014	Incidence is defined as the number of new infections during 2014.
HIV incidence per 100,000 population	CDC/Division of STD Prevention, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention	2014	Incidence is defined as the number of new infections during 2014.
HIV/AIDS hospitalization rate per 100,000 population	NA	NA	Data not available.
Syphilis incidence per 100,000 population	CDC/Division of STD Prevention, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention	2014	Incidence is defined as the number of new infections during 2014.
Intentional Injury			
Domestic assaults reports to police per 100,000 population	NA	NA	All offenses of assault between family or household members are reported as domestic assault.
Firearm deaths per 100,000 population	CDC/National Center for Health Statistics, National Vital Statistics System	2013	ICD-10 W32-W34 ,X72-X74, X93-X95, Y22-Y24, Y350 or U014.
Intentional self-injury (Youth)	NA	NA	Data not available.
Lifetime rape/non-consensual sex (among females)	NA	NA	Data not available.
Nonfatal child maltreatment per 1,000 population	U.S. Department of Health & Human Services, Administration for Children and Families, Administration on Children, Youth and Families, Children's Bureau	2011	Rates are unique child victims per 1,000 population under age 18. U.S. rate for 2011.
Reported rape per 100,000 population	FIB Uniform Crime Reports	2013	Includes rape by force and attempted forcible rape. Excludes carnal abuse without force (statutory rape) and other sex offenses.

Maine Shared Community Health Needs Assessment: 2015

U.S. Data Sources

Indicator	Data Source	Year(s)	Other Notes
Suicide deaths per 100,000 population	CDC/National Center for Health Statistics, National Vital Statistics System	2013	ICD-10 U03 X60-X84 or Y87.0
Violence by current or former intimate partners in past 12 months (among females)	NA	NA	Data not available.
Violent crime rate per 100,000 population	FIB Uniform Crime Reports	2013	Reported violent crime offenses. Violent crime includes murder, rape, robbery and aggravated assault.
Unintentional Injury			
Always wear seatbelt (adults)	NA	NA	Data not available.
Always wear seatbelt (high school students)	YRBS	2013	High School students who report they always wear a seatbelt when riding in a vehicle.
Traumatic brain injury related emergency department visits (all intents) per 10,000 population	National Center for Health Statistics, Health Indicators Warehouse	2010	ICD 9 CM 800–909.2, 909.4, 909.9–994.9, 995.5–995.59 or 995.80–995.85 or any external cause of injury code is ICD 9 CM E800–E869, E880–E929 or E950–E999, and the principal or any other diagnosis is ICD-9-CM 800.00–801.99, 803.00–804.99, 850.0–850.9, 851.00–854.19, 950.1–950.3, 959.01 or 995.55.
Unintentional and undetermined intent poisoning deaths per 100,000 population	CDC/National Center for Health Statistics, National Vital Statistics System	2013	If you want the complete definition, it's "deaths of U.S. residents for which the underlying cause of death is ICD-10 X40-X49 or Y10-Y19.
Unintentional fall related deaths per 100,000 population	CDC/National Center for Health Statistics, National Vital Statistics System	2013	Deaths of U.S. residents for which the underlying cause of death is ICD-10 W00-W19.
Unintentional fall related injury emergency department visits per 10,000 population	NA	NA	Data not available.
Unintentional motor vehicle traffic crash related deaths per 100,000 population	CDC/National Center for Health Statistics, National Vital Statistics System	2013	Deaths of U.S. residents for which the underlying cause of death is ICD-10 V02-V04 (.1, .9), V09.2, V12-V14 (.3-.9), V19 (.4-.6), V20-V28 (.3-.9), V29 (.4-.9), V30-V39 (.4-.9), V40-V49 (.4-.9), V50-V59 (.4-.9), V60-V69 (.4-.9), V70-V79 (.4-.9), V80 (.3-.5), V81.1, V82.1, V83-V86 (.0-.3), V87 (.0-.8) or V89.2."
Occupational Health			
Deaths from work-related injuries (number)	U.S. Dept. of Labor	2013	Includes self-employed workers, owners of unincorporated businesses and farms, paid and unpaid family workers, members of partnerships and may include owners of incorporated businesses.
Nonfatal occupational injuries (number)	NA	NA	Data not available.
Mental Health			
Adults who have ever had anxiety	NA	NA	Data not available.
Adults who have ever had depression	BRFSS	2013	Adults who have ever been told by a doctor or other healthcare provider that they have a depressive disorder.
Adults with current symptoms of depression	NA	NA	Data not available.
Adults currently receiving outpatient mental health treatment	NA	NA	Data not available.
Co-morbidity for persons with mental illness	NA	NA	Data not available.
Mental health emergency department rates per 100,000 population	NA	NA	Data not available.

Maine Shared Community Health Needs Assessment: 2015

U.S. Data Sources

Indicator	Data Source	Year(s)	Other Notes
Sad/hopeless for two weeks in a row (High School Students)	YRBS	2013	During the past 12 months, did you ever feel so sad or hopeless almost every day for two weeks or more in a row that you stopped doing some usual activities? Percentage of students who answered "Yes".
Seriously considered suicide (High School Students)	YRBS	2013	During the past 12 months, did you ever seriously consider attempting suicide? Percentage of students who answered "Yes".
Physical Activity, Nutrition and Weight			
Fewer than two hours combined screen time (High School Students)	NA	NA	Data not available.
Fruit and vegetable consumption (High School Students)	NA	NA	Data not available.
Fruit consumption among Adults 18+ (less than one serving per day)	BRFSS	2013	Adults with less than one serving per day of fruits or fruit juice.
Met physical activity recommendations (Adults)	BRFSS	2013	Adults who reported doing enough physical activity to meet the aerobic and strengthening recommendations.
Physical activity for at least 60 minutes per day on five of the past seven days (High School Students)	YRBS	2013	Percentage of students who were physically active for a total of at least 60 minutes per day on five of the past seven days.
Sedentary lifestyle – no leisure-time physical activity in past month (Adults)	BRFSS	2013	Adults reporting that during the past month, other than their regular job, they did not participate in any physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise.
Soda/sports drink consumption (High School Students)	YRBS	2013	Percentage of students who drank at least one can, bottle, or glass of soda, sports drink, energy drink, or other sugar-sweetened beverage such as Gatorade, Red Bull, lemonade, sweetened tea or coffee drinks, flavored milk, Snapple, or Sunny Delight? (Not counting diet soda, other diet drinks, or 100% fruit juice.) per day during the past week.
Vegetable consumption among Adults 18+ (less than one serving per day)	BRFSS	2013	Adults with less than one serving per day of vegetables.
Obesity (Adults)	BRFSS	2013	Adults with a BMI of 30 or more.
Obesity (High School Students)	YRBS	2013	Percentage of students who were obese (i.e., at or above the 95th percentile for body mass index, by age and sex) -- SELF-REPORTED HEIGHT/WEIGHT.
Overweight (Adults)	BRFSS	2013	Adults with a BMI between 25.0 and 29.9.
Overweight (High School Students)	YRBS	2013	Percentage of students who were overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex) -- SELF-REPORTED HEIGHT/WEIGHT.
Pregnancy and Birth Outcomes			
Children with special health care needs	National Survey of Children with Special Health Care Needs	2009-2010	Survey respondents who reported that their child has a special health care need.
Infant deaths per 1,000 live births	CIA World Factbook	2012	Number of babies who died before their first birthday per 1,000 live births. Average annual number of infant deaths and infant mortality rate might be slightly underestimated due to possible missing out-of-state deaths of U.S. infants in 2010.
Live births for which the mother received early and adequate prenatal care	CDC/National Center for Health Statistics, National Vital Statistics System	2013	Defined as an adequate or adequate-plus rating on the Kotelchuck Adequacy of Prenatal Care Utilization Index.

Maine Shared Community Health Needs Assessment: 2015

U.S. Data Sources

Indicator	Data Source	Year(s)	Other Notes
Live births to 15-19 year olds per 1,000 population	CDC/National Center for Health Statistics, National Vital Statistics System	2013	Defined as the number of live births among 15- to 19-year-old U.S. women per 1,000 population.
Low birth weight (<2500 grams)	U.S. Department of Health and Human Services, Health Resources and Services Administration, Maternal and Child Health Bureau	2013	Low birth weight defined as less than 2500 grams.
Substance and Alcohol Abuse			
Alcohol-induced mortality per 100,000 population	CDC/National Center for Health Statistics, National Vital Statistics System	2013	ICD-10 - E24.4 , F10, G31.2, G62.1, G72.1, I42.6, K29.2, K70, K85.2, K86.0, R78.0, X45, X65 or Y15
Binge drinking of alcoholic beverages (High School Students)	YRBS	2013	During the past 30 days, on how many days did you have 5 or more drinks of alcohol in a row, that is, within a couple of hours? Percentage of students who answered at least 1 day.
Binge drinking of alcoholic beverages (Adults)	BRFSS	2013	Risk factor for binge drinking where binge drinking is defined as having 5 or more drinks on 1 occasion for men and 4 or more drinks on 1 occasion for women.
Chronic heavy drinking (Adults)	BRFSS	2013	At risk for heavy alcohol consumption (greater than two drinks per day for men and greater than one drink per day for women).
Drug-affected baby referrals received as a percentage of all live births	NA	NA	Data not available.
Drug-induced mortality per 100,000 population	CDC/National Center for Health Statistics, National Vital Statistics System	2013	The population figures for year 2013 are bridged-race estimates of the July 1 resident population, from the Vintage 2013 postcensal series released by NCHS on June 26, 2014.
Emergency medical service overdose response per 100,000 population	NA	NA	Data not available.
Opiate poisoning (ED visits) per 100,000 population	NA	NA	Data not available.
Opiate poisoning (hospitalizations) per 100,000 population	NA	NA	Data not available.
Past-30-day alcohol use (High School Students)	YRBS	2013	During the past 30 days, on how many days did you have at least one drink of alcohol? Percentage of students who answered at least 1 day.
Past-30-day inhalant use (High School Students)	NA	NA	Data not available.
Past-30-day marijuana use (Adults)	NA	NA	Data not available.
Past-30-day marijuana use (High School Students)	YRBS	2013	During the past 30 days, how many times did you use marijuana? Percentage of students who answered at least 1 time.
Past-30-day nonmedical use of prescription drugs (Adult)	NA	NA	Data not available.
Past-30-day nonmedical use of prescription drugs (High School Students)	NA	NA	Data not available.
Prescription Monitoring Program opioid prescriptions (days supply/pop)	NA	NA	Data not available.
Substance-abuse hospital admissions per 100,000 population	NA	NA	Data not available.
Tobacco Use			
Current smoking (Adults)	BRFSS	2013	Adults that reported having smoked at least 100 cigarettes in their lifetime and currently smoke.

Maine Shared Community Health Needs Assessment: 2015

U.S. Data Sources

Indicator	Data Source	Year(s)	Other Notes
Current smoking (High School Students)	YRBS	2013	During the past 30 days, on how many days did you smoke cigarettes? Percentage of students who answered at least 1 day.
Current tobacco use (High School Students)	YRBS	2013	Percentage of students who smoked cigarettes or cigars or used chewing tobacco, snuff, or dip on one or more of the past 30 days. (Note: Reports read "Percentage of students who smoked cigarettes and/or cigars and/or used chewing tobacco, snuff, or dip on one or more of the past 30 days").
Secondhand smoke exposure (Youth)	NA	NA	Data not available.