

V. Miscellaneous Medical Formulas

Medical formulas are available for infants, children and women who have been assessed to have a medical condition which precludes or restricts the use of conventional foods and necessitates the use of a formula. ***The medical foods are not to be authorized solely for the purpose of enhancing nutrient intake or managing body weight if the participant is able to eat conventional foods adequately.***

The formula list below is not all inclusive. Other formulas, such as metabolic formulas, may be provided on a case-by-case basis. Provision of medical formulas will be coordinated with MaineCare.

For WIC participants who are enrolled in the MaineCare program:

- ***Hypercaloric and hydrolysate formulas*** will be provided by WIC up to the federal maximum issuance per month. Any medically necessary amounts needed over WIC's federal maximum will be provided by MaineCare.
- All ***elemental (amino acid-based) formulas*** will be provided by MaineCare

If WIC participants who are not enrolled in MaineCare:

- The Maine CDC WIC Nutrition Program will provide the formula in amounts not to exceed the federal maximum issuance per month.

Prescriptions for these formulas must include:

- **specific product requested**
- **diagnosis**
- **length of time the product is needed**
- **amount prescribed per day**

Health care providers may use the ***Maine CDC WIC Nutrition Program/MaineCare Request/Prior Authorization for Medical Formula/WIC-Eligible Nutritionals*** form to obtain approval and prior authorization for MaineCare coverage. **This form must be sent to the local WIC office in order for MaineCare to approve coverage of the formula prescribed.**

Miscellaneous Medical Formulas																									
Product Name	Description	Packaging	Composition	Indications	Maximum Monthly WIC Issuance Amounts																				
Similac® PM 60/40 (Abbott Nutrition)	Low-iron, lower mineral content medical infant formula Gluten free Kosher Halal	Powder: 14.1 oz. (400g) can; 6 cans/case Reconstitution/can: 102 fl. oz. NDC: 70074-0608-50	CHO: lactose PRO: whey protein concentrate, sodium caseinate FAT: high oleic safflower, soy and coconut oils Contains vitamins and reduced mineral content Additional iron should be supplied from other sources. Standard dilution: 20 cal./fl. oz. Osmolality = 280 mOsm/kg H ₂ O	For infant requiring lower mineral intake, including those with impaired renal function. Calcium to phosphorous ratios designed to manage serum calcium disorders, such as hypercalcemia and hypocalcemia due to hyperphosphatemia. Conditions where infant is losing abnormal quantities of one or more electrolytes, it may be necessary to supply electrolytes from sources other than the formula. Low birth weight babies (<1500g at birth) may need additional calcium, phosphorus and sodium during periods of rapid growth	<table border="1"> <thead> <tr> <th>Non-breastfed infants</th> <th>0-3 months</th> <th>4-5 months</th> <th>6-11 months</th> </tr> </thead> <tbody> <tr> <td>Powder: 14.1 oz.</td> <td>8 cans</td> <td>9 cans</td> <td>6 cans</td> </tr> <tr> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <th>Partially breastfed infants</th> <th>1-3 months</th> <th>4-5 months</th> <th>6-11 months</th> </tr> <tr> <td>Powder: 14.1 oz.</td> <td>4 cans</td> <td>5 cans</td> <td>3 cans</td> </tr> </tbody> </table>	Non-breastfed infants	0-3 months	4-5 months	6-11 months	Powder: 14.1 oz.	8 cans	9 cans	6 cans					Partially breastfed infants	1-3 months	4-5 months	6-11 months	Powder: 14.1 oz.	4 cans	5 cans	3 cans
					Non-breastfed infants	0-3 months	4-5 months	6-11 months																	
Powder: 14.1 oz.	8 cans	9 cans	6 cans																						
Partially breastfed infants	1-3 months	4-5 months	6-11 months																						
Powder: 14.1 oz.	4 cans	5 cans	3 cans																						
Enfaport (Mead Johnson Nutrition)	Nutritionally complete, iron-fortified, high protein formula for infants with chylothorax or LCHAD deficiency 60:40 whey-to-casein ratio 84% of fat from MCT oil	RTU: 6 fl. oz. bottle, 24 bottles/case NDC: 0087-510525	CHO: corn syrup solids, lactose PRO: skim milk, whey protein concentrate FAT: MCT, soy, M. alpina and C. cohnii oils Contains vitamins and minerals Ready to use: 30 cal./fl. oz. Dilution instructions available for mixing to 20 -28 kcal/oz if needed Osmolality = 360 mOsm/kg H ₂ O	Enfaport is designed to meet the unique nutritional needs of infants with Chylothorax or LCHAD deficiency. Enfaport balances high levels of MCT oil for easier absorption, and has DHA and ARA, important fatty acids for infant development.	<table border="1"> <thead> <tr> <th>Non-breastfed infants</th> <th>0-3 months</th> <th>4-5 months</th> <th>6-11 months</th> </tr> </thead> <tbody> <tr> <td>RTU</td> <td>138 bottles</td> <td>152 bottles</td> <td>107 bottles</td> </tr> <tr> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <th>Partially breastfed infants</th> <th>1-3 months</th> <th>4-5 months</th> <th>6-11 months</th> </tr> <tr> <td>RTU</td> <td>64 bottles</td> <td>79 bottles</td> <td>56 bottles</td> </tr> </tbody> </table>	Non-breastfed infants	0-3 months	4-5 months	6-11 months	RTU	138 bottles	152 bottles	107 bottles					Partially breastfed infants	1-3 months	4-5 months	6-11 months	RTU	64 bottles	79 bottles	56 bottles
					Non-breastfed infants	0-3 months	4-5 months	6-11 months																	
RTU	138 bottles	152 bottles	107 bottles																						
Partially breastfed infants	1-3 months	4-5 months	6-11 months																						
RTU	64 bottles	79 bottles	56 bottles																						

Miscellaneous Medical Formulas																									
Product Name	Description	Packaging	Composition	Indications	Maximum Monthly WIC Issuance Amounts																				
3232 A (Mead Johnson Nutrition)	Nutritionally incomplete protein hydrolysate formula base that is to be used with added carbohydrate. Lactose Free Galactose Free 82% of fat from MCT oil	Powder: 16 oz cans, 6 cans/case Reconstitution/can: varies, according to mixing instructions NDC: 00087-042541	CHO: modified tapioca starch (serves as stabilizer) PRO: casein hydrolysate (from milk), L-amino acids FAT: MCT, corn oils Without added carbohydrate: 12.7 calories/ fl. oz. Calories/ounce and osmolality depend on mixing instructions and carbohydrates added	For dietary management of infants and children with disaccharidase deficiencies or other disorders of carbohydrate metabolism. Additional carbohydrate is needed. If used long-term, essential fatty acids should be considered.	<table border="1"> <thead> <tr> <th colspan="4">Maximum Monthly WIC Issuance Amounts</th> </tr> <tr> <th>Non-breastfed infants</th> <th>0-3 months</th> <th>4-5 months</th> <th>6-11 months</th> </tr> </thead> <tbody> <tr> <td>Powder: 16 oz.</td> <td>Dependent on Rx</td> <td>Dependent on Rx</td> <td>Dependent on Rx</td> </tr> <tr> <th>Partially breastfed infants</th> <th>1-3 months</th> <th>4-5 months</th> <th>6-11 months</th> </tr> <tr> <td>Powder: 16 oz.</td> <td>Dependent on Rx</td> <td>Dependent on Rx</td> <td>Dependent on Rx</td> </tr> </tbody> </table>	Maximum Monthly WIC Issuance Amounts				Non-breastfed infants	0-3 months	4-5 months	6-11 months	Powder: 16 oz.	Dependent on Rx	Dependent on Rx	Dependent on Rx	Partially breastfed infants	1-3 months	4-5 months	6-11 months	Powder: 16 oz.	Dependent on Rx	Dependent on Rx	Dependent on Rx
					Maximum Monthly WIC Issuance Amounts																				
Non-breastfed infants	0-3 months	4-5 months	6-11 months																						
Powder: 16 oz.	Dependent on Rx	Dependent on Rx	Dependent on Rx																						
Partially breastfed infants	1-3 months	4-5 months	6-11 months																						
Powder: 16 oz.	Dependent on Rx	Dependent on Rx	Dependent on Rx																						
Portagen® (Mead Johnson Nutritionals)	Nutritionally incomplete iron fortified formula with easily digestible fats. Suitable for lactose intolerance Low Residue Kosher Not recommended for infants 87% of fat from MCTs	Powder: 410g can, 6 cans/case Reconstitution/can: 64 fl. oz. NDC: 0087-511728	CHO: corn syrup solids, sugar PRO: sodium caseinate (from milk) FAT: MCT and corn oils Contains vitamins and minerals Standard dilution: 30 cal. /fl. oz. Osmolality = 350 mOsm/kg H ₂ O	For children and adults with defects in the intraluminal hydrolysis of fat (decreased pancreatic lipase, decreased bile salts), defective mucosal fat absorption (decreased mucosal permeability and/or absorptive surface) and/or defective lymphatic transport of fat (intestinal lymphatic obstruction) If used long term, supplementation of essential fatty acids and ultra-trace minerals should be considered. Not recommended for use as an infant formula.	<table border="1"> <thead> <tr> <th colspan="2">Children</th> </tr> </thead> <tbody> <tr> <td>Powder 16 oz.</td> <td>14 cans</td> </tr> </tbody> </table>	Children		Powder 16 oz.	14 cans																
Children																									
Powder 16 oz.	14 cans																								