

MDS 3.0 RUG III ADL Scoring Chart

The ADL index is used to split many of the Clinical Indicator categories. Composed from these ADL activities:

Bed Mobility – Items G011A1 and G0110A2

Transfer – Items G0110B1 and G0110B2

Eating – Items G0110G1, K0500B, K0700A, and K0700B

Toilet Use – Items G0110I1 and G0110I2

The ADL index score is determined as follows:

ADL Function		ADL Score
Bed mobility, Toilet use and Transfer:		
	Self-Performance (column 1) is coded on the MDS 3.0 as:	
	Independence or Supervision (item coded 0,1, or 7)	1
	Limited Assistance (item coded 2)	3
	Extensive Assistance or Total Dependence (item coded 3,4 or 8) AND Support Provided (Column 2) is:	
	<ul style="list-style-type: none"> None, set-up only or 1 person physical assist (item coded 0, 1, or 2) 	4
<ul style="list-style-type: none"> 2+ persons physical assist or activity did not occur (item coded 3 or 8*) 	5	
Eating:		
IF:	Resident receives Parenteral/IV (K0500A = 1)	3
	Tube Feeding (K0500B = 1) AND one of the following:	3
	<ul style="list-style-type: none"> 51% or more total calories from parenteral or tube feeding (K0700A = 3) 	
	<ul style="list-style-type: none"> 26-50% total calories from parenteral or tube feeding (K0700A = 2) AND fluid intake is 501+ ml per day (K0700B = 2) 	
IF NOT:	Self-Performance (column 1) is:	
	Independent or Supervision (item coded 0, 1, 7)	1
	Limited Assistance (item coded 2)	2
	Extensive Assistance or Total Dependence or Activity Did Not Occur (item coded 3, 4, or 8)	3

The scores for each ADL variable are added to compute the ADL index score. The ADL Index score will range from 4 to 18 for each resident.