

# tick ID



KNOW THEM. PREVENT THEM.



## Deer Tick (Black-Legged Tick)



nymph



adult male



adult female

- Deer ticks may transmit the agents that cause Lyme disease, anaplasmosis, and babesiosis
- **What bites:** nymphs and adult females
- **When:** anytime temperatures are above freezing, greatest risk is spring through fall

ACTUAL SIZE



nymph  
(1/32"–1/16")



adult  
(1/8")



engorged adult  
(up to 1/2")

- Dog ticks do not transmit the agent that causes Lyme disease
- **What bites:** adult females
- **When:** April–August through fall

## Dog Tick



adult male



adult female

*(examples are not actual size, dog tick nymphs are rarely found on humans or their pets)*

## prevent the bite

- Wear light-colored protective clothing
- Use EPA-approved insect repellent on skin or clothing
- Use caution in tick infested areas
- Perform daily tick checks
- Protect your pets, use repellents, acaricides, and a Lyme disease vaccine for dogs



## tick removal

Remove ticks immediately. They usually need to attach for 24 hours to transmit Lyme disease. Consult a physician if you remove an engorged deer tick.

### Using a tick spoon:

- Place the wide part of the notch on the skin near the tick (hold skin taut if necessary)
- Applying slight pressure downward on the skin, slide the remover forward so the small part of the notch is framing the tick
- Continuous sliding motion of the remover detaches the tick

### Using tweezers:

- Grasp the tick close to the skin with tweezers
- Pull gently until the tick lets go